

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
21 December 2000 (21.12.2000)

PCT

(10) International Publication Number
WO 00/76982 A1

- (51) International Patent Classification⁷: C07D 215/44, 219/12, A61K 31/47, A61P 37/06
- (74) Agent: SHISHIMA, Gina, N.; Fulbright & Jaworski, L.L.P., Suite 2400, 600 Congress Avenue, Austin, TX 78701 (US).
- (21) International Application Number: PCT/US00/16723
- (22) International Filing Date: 16 June 2000 (16.06.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/139,544 16 June 1999 (16.06.1999) US
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant: UNIVERSITY OF IOWA RESEARCH FOUNDATION [US/US]; 100 Oakdale Campus #214-TIC, Iowa City, IA 52242-5000 (US).
- (72) Inventors: MACFARLANE, Donald, E.; 620 S. Summit Street, Iowa City, IA 52240 (US). STREKOWSKI, Lucjan; 6091 Windsong Way, Stone Mountain, GA 30087 (US). MANZEL, Lori; 3316 Timber Valley Ct. NE, Cedar Rapids, IA 52402 (US). ISMAIL, Fyaz; Department of Physical Sciences, University of Hertfordshire, Hatfield AL10 9AB, Hertfordshire (GB). BARLIN, Gordon, B.; John Curtin School of Medical Research, Room D3044A, The Australian National University, Canberra, ACT 0200 (AU).
- Published:**
— With international search report.
— Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: ANTAGONISM OF IMMUNOSTIMULATORY CPG-OLIGONUCLEOTIDES BY 4-AMINOQUINOLINES AND OTHER WEAK BASES

(57) Abstract: The present invention relates generally to the field of immunology. More particularly it concerns compositions and methods for inhibiting stimulation of the immune system. The compounds and methods comprise compounds that are analogs and derivatives of chloroquine, such as 4-aminoquinolines, and other weak bases. They can be used in preventative and therapeutic treatments of autoimmune diseases and phenomena, transplant rejection such as host-versus-graft disease, and sepsis.

WO 00/76982 A1

DESCRIPTION

ANTAGONISM OF IMMUNOSTIMULATORY CPG-OLIGONUCLEOTIDES BY 4-AMINOQUINOLINES AND OTHER WEAK BASES

BACKGROUND OF THE INVENTION

The present application claims the benefit of U.S. Provisional Application Serial Number 60/139,544 filed on June 16, 1999. The entire text of the above-referenced disclosure is herein incorporated by reference.

1. Field of the Invention

The present invention relates generally to the field of immunology. More particularly, it concerns preventative and therapeutic compositions and methods that inhibit stimulation of the immune system.

2. Description of Related Art

While stimulation of the immune systems prevents and controls infection, it can have an adverse physiological effect, as is the case with autoimmune diseases and phenomena, with rejection of cells and tissues during adoptive immunotherapy and transplants, and with invasions by pathogens. Inhibition of this stimulation can have beneficial therapeutic results. However, new and more effective treatments to effect inhibition of immunostimulation are still needed.

Autoimmunity is generally caused by aberrations in lymphocyte activities. While the precise cause of autoimmunity is not known, it most likely involves a mechanistic failure in at least one of the steps of maintaining self-tolerance to the body's own antigens. Several factors are thought to play a part in the development of autoimmunity, including the host's genetic makeup. Autoimmune diseases afflict approximately 1% to 2% of the human population. Autoimmune phenomena, are the result of a disease, for example myocardial infarction, that may cause damage to tissue, which consequently effects the release of immunogenic tissue antigens; this

condition, unlike autoimmune disease, is unrelated to the pathogenesis of the disease that caused it.

There are a wide variety of autoimmune diseases. They are classified as either organ-specific based on the primary site of injury or systemic (see Table 1). There are three ways in which damage or injury to tissue is caused by autoimmune disease: cell-mediated immunity, cell lysis and autoantibody-induced release of inflammatory mediators, and immune complex disease.

Cell-mediated immunity occurs when sensitized T cells directly damage cells or release lymphokines that augment the inflammatory reaction. An association of an autoantibody with its antigen in intercellular fluid causes cell lysis and autoantibody-induced release of inflammatory mediators. This interaction results in release of inflammatory mediators, induction of the complement pathway, or activation of cytotoxic cells, which can trigger cell lysis. The third mechanism, immune complex disease, involves a reaction between circulating autoantibodies and antigens on the cell surface. This complex becomes deposited in tissue such as the joints, blood vessels, and glomeruli, causing complement to be fixed and subsequent inflammation and tissue damage.

Rejection of cells and tissue can involve rejection of the graft by the host. The body's own cells are identified as self because of a complex series of cell surface molecules known as major histocompatibility molecules (MHC). Rejection of cells and tissues can occur following transplantation of cells or organs or after adoptive immunotherapy has been implemented. In graft-versus host disease (GVHD), the grafted immune system attacks the host cells. One example in which GVHD becomes particularly significant is bone marrow transplantation (BMT), which is frequently used for the treatment of a variety of bone marrow-related disorders and in cancer therapy to replace bone marrow cells lost to chemotherapy and radiation treatment. In severe cases of GVHD, a patient's compromised immune system gives rise to many complications including those in the liver, causing jaundice, in the skin causing rash,

and in the gastrointestinal tract, including diarrhea, anorexia, nausea and vomiting, malabsorption, abdominal pain ileus, and ascites formation.


Sepsis is the primary cause of death in the intensive care unit with more than 400,000 cases in the United States annually. It can be caused by infection by a pathogen, such as viruses, bacteria, fungi, and parasites, which triggers host defenses. This may result in activation of innate immunity, particularly, an inflammatory response, which consequently promotes deleterious effects (collectively termed "sepsis") including shock, respiratory distress, capillary leaks, renal failure, jaundice, bleeding, coma and death.

Despite this information, preventative and therapeutic treatment to inhibit stimulation of the immune system is still needed. Oligodeoxynucleotides (ODN), bacterial DNA, and phosphorothioate oligodeoxynucleotides with unmethylated CpG-motifs are immunostimulatory and may contribute to autoimmunity. Thus, they can serve as a model system to identify compounds and methods that effect inhibition of immunostimulation. Using CpG-ODN, a number of compounds have been identified as possessing this property. For example, a number of quinoline derivatives that are active against stimulation by CpG-ODN have been shown to induce remission of rheumatoid arthritis and lupus erythematosus (Fox, 1993; Wallace, 1994). Chloroquine and a number of structural analogs specifically and powerfully inhibit this effect at nanomolar concentration. Therefore, inhibition of CpG-ODN immunostimulation can be effected generally in the treatment of autoimmune diseases and phenomena, sepsis, and transplantation rejection, including graft-versus-host disease.

SUMMARY OF THE INVENTION

The inhibition of immune stimulation in cases such as autoimmune diseases, tissue transplantation and the like would be therapeutically beneficial. Currently, methods to inhibit stimulation of the immune system are limited. Therefore, it is a goal of the present invention to provide methods and compositions for inhibiting immunostimulation in a subject.

In one embodiment of the invention, a method of inhibiting immunostimulation in a subject by administering an effective amount of a substituted 4-quinolinamine composition to the subject comprising a compound having the structural formula A is provided.




(A)

A substituted 4-quinolinamine composition of formula A, comprises groups R_A , R_B , R_2 , R_3 , R_5 , R_6 , R_7 and R_8 , wherein R_A is hydrogen atom or a lower alkyl group, R_B is a substituted or unsubstituted alkyl, alkenyl or alkynyl secondary or tertiary amine, R_2 is a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, an substituted or unsubstituted anthracyl group, a substituted or unsubstituted phenanthryl group or a substituted or unsubstituted styryl group, R_3 is a hydrogen atom, R_5 is a hydrogen atom, R_6 is a hydrogen atom or a halogen atom, R_7 is a hydrogen atom or a halogen atom, R_8 is a hydrogen atom and pharmaceutically acceptable salts thereof. In preferred embodiments, the R_2 phenyl, naphthyl, anthracyl, styryl or phenanthryl group substitution further comprise one or more substitutions selected from the group consisting of an alkyl group, an alkoxy group, an alkoxyalkyl group, an ester group, an alkylamino group, a dialkylamino group, a cyclic amino group, a halogen atom, and any combination thereof. In some aspects of the invention, the R_2 substitution is an *N*-substituted aminomethyl group, a cyclic aminomethyl group, or an aminomethyl group, additionally substituted at the nitrogen atom with a cyclic aminoalkyl group. In particularly preferred embodiments, a cyclic amino group is a piperazino group, a piperidino group, a pyrrolidino group, an

imidazolyl group, a pyridyl, or a morpholino group. In another embodiment, the R_B substituted alkyl is selected from the group consisting of a cyclic amino group, furyl, thienyl, phenyl, alkylamino group, dialkylamino group, and any combination thereof. It is contemplated that phenyl includes a substituted phenyl. In preferred embodiments, a cyclic amino group is a piperazino group, a piperidino group, a pyrrolidino group, an imidazolyl group, a pyridyl group, or a morpholino group. The cyclic amino groups in R_B or R_2 may be further substituted with an alkyl group, for example, an *N*-methylpiperazino or an *N*-pyrrolidinoalkyl. It is further contemplated that there is a covalent bond formed between R_A and R_B , for example, to form an *N*-methylpiperazino.

In another embodiment, the present invention provides a method of inhibiting immunostimulation in a subject comprising administering an effective amount of a substituted 4-quinolinamine composition to said subject comprising a compound having the structural formula B.




(B)

A substituted 4-quinolinamine composition of formula B, comprises a phenyl group, R_2 , R_3 , R_5 , R_6 , R_7 and R_8 , wherein R_2 is a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted anthracyl group, a substituted or unsubstituted styryl group or a substituted or unsubstituted phenanthryl group, R_3 is a hydrogen atom, R_5 is a hydrogen atom, R_6 is

a hydrogen atom or a halogen atom, R_7 is a hydrogen atom, a halogen atom or an alkyl halogen atom, R_8 is a hydrogen atom and pharmaceutically acceptable salts thereof. In a preferred embodiment, when the R_2 phenyl, naphthyl, anthracyl, styryl or phenanthryl group comprises substitutions, these are defined as one or more substitutions selected from the group consisting of an alkyl group, an alkoxy group, an alkoxyalkyl group, an ester group, an alkylamino group, a dialkylamino group, a cyclic amino group, a halogen atom or any combination thereof. In particularly preferred embodiments, a cyclic amino group is a piperazino group, a piperidino group, a pyrrolidino group, an imidazolino group or a morpholino group.

The phenyl group additionally may be substituted at positions R_9 , R_{10} , R_{11} , R_{12} and R_{13} , wherein the substituted phenyl group comprises one or more substitutions selected from the group consisting of an alkyl group, an alkoxy group, an alkoxyalkyl group, an ester group, an alkylamino group, a dialkylamino group, a cyclic amino group, a furan group, a thiophene group, a halogen atom or any combination thereof. In preferred embodiments, a cyclic amino group is a piperazino group, a piperidino group, a pyrrolidino group, an imidazolino group or a morpholino group.

In further embodiments of the invention, a method of inhibiting immunostimulation in a subject comprises administering an effective amount of a substituted 4-quinolinamine composition to the subject comprising a compound having the structural formula C.




(C)

A substituted 4-quinolinamine composition of formula C, comprises a phenyl group spatially distanced from the quinolinamine by an alkyl group of 0 to 4 CH₂ molecules, R₂, R₃, R₅, R₆, R₇ and R₈, wherein R₂ is a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted anthracyl group, a substituted or unsubstituted styryl group or a substituted or unsubstituted phenanthryl group, R₃ is a hydrogen atom, R₅ is a hydrogen atom, R₆ is a hydrogen atom or a halogen atom, R₇ is a hydrogen atom, a halogen atom or an alkyl halogen atom, R₈ is a hydrogen atom and pharmaceutically acceptable salts thereof. In preferred embodiments, the R₂ phenyl, naphthyl, anthracyl, styryl or phenanthryl group substitution comprises one or more substitutions selected from the group consisting of an alkyl group, an alkoxy group, an alkoxyalkyl group, an ester group, an alkylamino group, an dialkylamino group, a cyclic amino group, a halogen atom or any combination thereof. In particularly preferred embodiments, a cyclic amino group is a piperazino group, a piperidino group, a pyrrolidino group, an imidazolino group or a morpholino group.

In other embodiments of the invention, the phenyl group additionally may be substituted at positions R₉, R₁₀, R₁₁, R₁₂ and R₁₃, wherein the substituted phenyl group comprises one or more substitutions selected from the group consisting of an alkyl group, an alkoxy group, an alkoxyalkyl group, an ester group, an alkylamino group, an dialkylamino group, a cyclic amino group, a furan group, a thiophene group, a halogen atom, or any combination thereof.

In another embodiment, a method of inhibiting immunostimulation in a subject comprises administering an effective amount of a substituted bis-4-quinolinamine composition to the subject comprising a compound having the structural formula D.




(D)

A substituted bis-4-quinolinamine composition having the structural formula D, comprises R_B on the first 4-quinolinamine covalently attached to $R_{B'}$ on the second 4-quinolinamine by linker group Y, R_2 is a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted anthracyl group, a substituted or unsubstituted styryl group or a substituted or unsubstituted phenanthryl group, R_3 is a hydrogen atom, R_5 is a hydrogen atom, R_6 is a hydrogen atom or a halogen atom, R_7 is a hydrogen atom or a halogen atom, R_8 is a hydrogen atom, $R_{2'}$ is a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, an substituted or unsubstituted anthracyl group, a substituted or unsubstituted styryl group or a substituted or unsubstituted phenanthryl group, $R_{3'}$ is a hydrogen atom, $R_{5'}$ is a hydrogen atom, $R_{6'}$ is a hydrogen atom or a halogen atom, $R_{7'}$ is a hydrogen atom or a halogen atom, $R_{8'}$ is a hydrogen atom and pharmaceutically acceptable salts thereof.

In other embodiments, the R_2 phenyl, naphthyl, anthracyl, styryl or phenanthryl group substitution is further defined as one or more substitutions selected from the group consisting of an alkyl group, an alkoxy group, an alkoxyalkyl group, an ester group, an alkylamino group, an dialkylamino group, a cyclic amino group, a halogen atom or any combination thereof and the $R_{2'}$ phenyl, naphthyl, anthracyl, styryl or phenanthryl group substitution is further defined as one or more substitutions selected from the group consisting of an alkyl group, an alkoxy group, an alkoxyalkyl group, an ester group, an alkylamino group, an dialkylamino group, a cyclic amino group, a halogen atom or any combination thereof.

In preferred embodiments, a cyclic amino group is a piperazino group, a piperidino group, a pyrrolidino group, an imidazolino group or a morpholino group. In particular embodiments, the linker group Y is an alkyl group, an ester group, an alkoxyalkyl group, an alkylamino group, a dialkylamino group, amido group, a cyclohexane, an piperazino group or any combination thereof.

In other embodiments of the present invention, a method of inhibiting immunostimulation in a subject comprising administering an effective amount of a substituted bis-9-aminoacridine composition to the subject, comprising a compound




having the structural formula E, is provided.

(E)

A substituted bis-9-aminoacridine of formula E, comprises R_B on the first 9-aminoacridine covalently attached to $R_{B'}$ on the second 9-aminoacridine by linker group Y. A substituted bis-9-aminoacridine of formula E further comprises OR_2 , OR_2' , R_3 , R_3' , X and X', wherein OR_2 is a lower alkyl group, OR_2' is a lower alkyl group, R_3 is a hydrogen atom or a lower alkoxy group, R_3' is a hydrogen atom or a lower alkoxy group, X is a halogen atom, X' is a halogen atom and pharmaceutically acceptable salts thereof. In other embodiments, the linker group Y is an alkyl group, an ester group, an alkoxyalkyl group, an alkylamino group, a dialkylamino group, amido group, a cyclohexane, an piperazino group or any combination thereof.


In yet another embodiment, a method of inhibiting immunostimulation in a subject comprises administering an effective amount of a composition to a subject comprising a compound having the structural formula F, wherein a 4-quinolinamine at position R_B is covalently linked to a 9-aminoacridine at position R_B , by linker group Y.



(F)

The 4-quinolinamine, 9-aminoacridine composition of formula F comprises R_A , R_B , R_2 , R_3 , R_5 , R_6 , R_7 , R_8 , R_B' , OR_2' , R_3' and X , wherein R_A is atom or a lower alkyl group, R_2 is a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, an substituted or unsubstituted anthracyl group, a substituted or unsubstituted styryl group or a substituted or unsubstituted phenanthryl group, R_3 is a hydrogen atom, R_5 is a hydrogen atom, R_6 is a hydrogen atom or a halogen atom, R_7 is a hydrogen atom or a halogen atom, R_8 is a hydrogen atom, OR_2' is a lower alkyl group, R_3' is a hydrogen atom, or a lower alkoxy group, X is a halogen atom and pharmaceutically acceptable salts thereof. In certain embodiments, the R_2 phenyl, naphthyl, anthracyl, styryl or phenanthryl group substitution is further defined as one or more substitutions selected from the group consisting of an alkyl group, an alkoxy group, an alkoxyalkyl group, an ester group, an alkylamino group, an dialkylamino group, a cyclic amino group, a halogen atom or any combination thereof. In preferred embodiments, a cyclic amino group is a piperazino group, a piperidino group, a pyrrolidino group, an imidazolino group or a morpholino group. In other embodiments, the Y linker is an alkyl group, an ester group, an alkoxyalkyl group, an alkylamino group, an dialkylamino group, amido group, a cyclohexane, an piperazino group or any combination thereof.


In other embodiments of the present invention, 4-quinolinamine and 9-aminoacridine compositions are provided. Thus, in one embodiment, a substituted 4-quinolinamine composition having the structural formula A is provided.



(A)

R_A is hydrogen atom or a lower alkyl group, R_B is a substituted or unsubstituted alkyl, alkenyl or alkynyl secondary or tertiary amine, R_2 is a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, an substituted or unsubstituted anthracyl group, a substituted or unsubstituted phenanthryl group or a substituted or unsubstituted styryl group, R_3 is a hydrogen atom, R_5 is a hydrogen atom, R_6 is a hydrogen atom or a halogen atom, R_7 is a hydrogen atom or a halogen atom, R_8 is a hydrogen atom and pharmaceutically acceptable salts thereof. In particular embodiments, the R_2 phenyl, naphthyl, anthracyl, styryl or phenanthryl group substitution is further defined as one or more substitutions selected from the group consisting of an alkyl group, an alkoxy group, an alkoxyalkyl group, an ester group, an alkylamino group, a dialkylamino group, a cyclic amino group, a halogen atom and any combination thereof. In preferred embodiments the cyclic amino group is a piperazino group, a piperidino group, a pyrrolidino group, an imidazolino group or a morpholino group. In yet other embodiments, the R_B alkyl substitution is selected from the group consisting of a cyclic amino group, furan and thiophene, wherein the cyclic amino group is a piperazino group, a piperidino group, a pyrrolidino group, an imidazolino group or a morpholino group.

In other embodiments of the present invention a substituted 4-quinolinamine composition is provided having the structural formula B.



(B)

5

The phenyl group can be substituted at R₉, R₁₀, R₁₁, R₁₂ and R₁₃, wherein a substituted phenyl group comprises one or more substitutions selected from the group consisting of an alkyl group, an alkylaminoalkyl group, an alkoxy group, an alkoxyalkyl group, an ester group, an alkylamino group, an dialkylamino group, a cyclic amino group, a furan group, a thiophene group, a halogen atom or any combination thereof. R₂ is a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted anthracyl group, a substituted or unsubstituted styryl group or a substituted or unsubstituted phenanthryl group, R₃ is a hydrogen atom, R₅ is a hydrogen atom, R₆ is a hydrogen atom or a halogen atom, R₇ is a hydrogen atom, a halogen atom or an alkyl halogen atom, R₈ is a hydrogen atom and pharmaceutically acceptable salts thereof. In other embodiments, a cyclic amino group is a piperazino group, a piperidino group, a pyrrolidino group, an imidazolino group or a morpholino group.


20

In yet other embodiments, the R₂ phenyl, naphthyl, anthracyl, styryl or phenanthryl group substitution is further defined as one or more substitutions selected from the group consisting of an alkyl group, an alkoxy group, a hydroxy group, an alkoxyalkyl group, an ester group, an alkylamino group, an dialkylamino group, a cyclic amino group, a halogen atom or any combination thereof. In certain

25

embodiments, a cyclic amino group is a piperazino group, a piperidino group, a pyrrolidino group, an imidazolino group or a morpholino group.

In particular embodiments, a substituted 4-quinolinamine composition having the structural formula C is provided.



(C)

5


The phenyl group can be unsubstituted or substituted at R₉, R₁₀, R₁₁, R₁₂ and R₁₃, wherein the substituted phenyl group is distanced spatially from the quinolinamine by an alkyl group of 0 to 4 CH₂ and comprises one or more
 10 substitutions selected from the group consisting of an alkyl group, an alkoxy group, an alkoxyalkyl group, an ester group, an alkylamino group, a dialkylamino group, a cyclic amino group, a furan group, a thiophene group, a halogen atom, or any combination thereof. R₂ is a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted anthracyl group, a
 15 substituted or unsubstituted styryl group or a substituted or unsubstituted phenanthryl group, R₃ is a hydrogen atom, R₅ is a hydrogen atom, R₆ is a hydrogen atom or a halogen atom, R₇ is a hydrogen atom, a halogen atom or an alkyl halogen atom, R₈ is a hydrogen atom and pharmaceutically acceptable salts thereof. In certain
 20 embodiments, a cyclic amino group is a piperazino group, a piperidino group, a pyrrolidino group, an imidazolino group or a morpholino group.

In other embodiments, the R₂ phenyl, naphthyl, anthracyl, styryl or phenanthryl group substitution is further defined as one or more substitutions selected from the group consisting of an alkyl group, an alkoxy group, an alkoxyalkyl group, an ester

group, an alkylamino group, an dialkylamino group, a cyclic amino group, a halogen atom or any combination thereof. In yet other embodiments, a cyclic amino group is a piperazino group, a piperidino group, a pyrrolidino group, an imidazolino group or a morpholino group.

5

In another embodiment of the invention, a substituted bis -4-quinolinamine composition having the structural formula D is provided.



(D)


10

The R_B on the first 4-quinolinamine is covalently attached to R_{B'} on the second 4-quinolinamine by linker group Y, wherein linker Y is an alkyl group, an ester group, an alkoxyalkyl group, an alkylaminoalkyl group, an alkylamino group, an dialkylamino group, amido group, a cyclohexane, an piperazino group or any combination thereof. R₂ is a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted anthracyl group, a substituted or unsubstituted styryl group or a substituted or unsubstituted phenanthryl group, R₃ is a hydrogen atom, R₅ is a hydrogen atom, R₆ is a hydrogen atom or a halogen atom, R₇ is a hydrogen atom or a halogen atom, R₈ is a hydrogen atom, R_{2'} is a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, an substituted or unsubstituted anthracyl group, a substituted or unsubstituted styryl group or a substituted or unsubstituted phenanthryl group, R_{3'} is a hydrogen atom, R_{5'} is a hydrogen atom, R_{6'} is a hydrogen atom or a halogen atom, R_{7'} is a hydrogen atom or a halogen atom, R_{8'} is a hydrogen atom and pharmaceutically acceptable salts thereof.

25

In certain embodiments, the R_2 phenyl, naphthyl, anthracyl, styryl or phenanthryl group substitution is further defined as one or more substitutions selected from the group consisting of an alkyl group, an alkoxy group, an alkoxyalkyl group, a hydroxy group, an ester group, an alkylamino group, an dialkylamino group, a cyclic amino group, a halogen atom or any combination thereof. In preferred embodiments, a cyclic amino group is a piperazino group, a piperidino group, a pyrrolidino group, an imidazolino group or a morpholino group. In other embodiments, the R_2 phenyl, naphthyl, anthracyl, styryl or phenanthryl group substitution is further defined as one or more substitutions selected from the group consisting of an alkyl group, an alkoxy group, a hydroxy group, an alkoxyalkyl group, an ester group, an alkylamino group, an dialkylamino group, a cyclic amino group, a halogen atom or any combination thereof. In particular embodiments, a cyclic amino group is a piperazino group, a piperidino group, a pyrrolidino group, an imidazolino group or a morpholino group.

In certain embodiments, a substituted bis-9-aminoacridine composition having the structural formula E is provided.



(E)


20

The R_B on the first 9-aminoacridine is covalently attached to $R_{B'}$ on the second 9-aminoacridine by linker group Y, wherein linker Y is an alkyl group, an ester group, an ether group, an alkylamino group, an dialkylamino group, amido group, a cyclohexane, an piperazino group or any combination thereof. OR_2 is a lower alkyl group, OR_2' is a lower alkyl group, R_3 is a hydrogen atom or a lower alkoxy group, R_3'

25

is a hydrogen atom or a lower alkoxy group, X is a halogen atom, X' is a halogen atom and pharmaceutically acceptable salts thereof.

In another embodiment, 4-quinoliamine, 9-aminoacridine composition having the structural formula F is provided.



(F)

The R_B position on the 4-quinoliamine is covalently attached to the R_B position of the 9-aminoacridine by linker Y, wherein linker Y is an alkyl group, an ester group, an alkoxyalkyl group, an alkylamino group, amido group, a cyclohexanedieyl, an piperazino group or any combination thereof. R_A is hydrogen atom or a lower alkyl group, R₂ is a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, an substituted or unsubstituted anthracyl group, a substituted or unsubstituted styryl group or a substituted or unsubstituted phenanthryl group, R₃ is a hydrogen atom, R₅ is a hydrogen atom, R₆ is a hydrogen atom or a halogen atom, R₇ is a hydrogen atom or a halogen atom, R₈ is a hydrogen atom, OR₂' is a lower alkyl group, R₃' is a hydrogen atom, or a lower alkoxy group, X is a halogen atom and pharmaceutically acceptable salts thereof.

20

In particular embodiments, the R₂ phenyl, naphthyl, anthracyl, styryl or phenanthryl group substitution is further defined as one or more substitutions selected from the group consisting of an alkyl group, an alkoxy group, an alkoxyalkyl group, an ester group, an alkylamino group, an dialkylamino group, a cyclic amino group, a halogen atom or any combination thereof. In certain embodiments, a cyclic amino group is a piperazino group, a piperidino group, a pyrrolidino group, an imidazolino group or a morpholino group.

25

It is also contemplated in the present invention that various other substitutions, such as hydroxy groups and various substituted alkyl groups can be made at the R_B and R₂ positions of the substituted 4-quinoliamine.

5

The term "lower alkyl" in the context of the present invention represents straight and branched alkyl groups up to 7 carbon atoms. As used herein the specification, "a" or "an" may mean one or more. As used herein in the claim(s), when used in conjunction with the word "comprising," the words "a" or "an" may mean one or more than one. As used herein "another" may mean at least a second or more.

10

Other objects, features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples, while indicating preferred embodiments of the invention, are given by way of illustration only, since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description.

15

20

BRIEF DESCRIPTION OF THE DRAWINGS

The following drawings form part of the present specification and are included to further demonstrate certain aspects of the present invention. The invention may be better understood by reference to one or more of these drawings in combination with the detailed description of specific embodiments presented herein.

25

FIG. 1A.-1P. Various analogs with some inhibitory activity against CpG ODN immunostimulation.

30

FIG. 2. Structures of analogs used. The numbers are those in the first column of Table 9.

FIG. 3. Uptake of analogs by cells. WEHI 231 cells were incubated with fluorescent analogs (0.1-20 μ M) for 30 min. The concentration of the analogs in the supernatant medium (free) and cellular pellet (bound) was then determined as described in Methods.

FIG. 4. Efflux of analog. WEHI 231 cells were incubated with 20 μ M analog for 30 min. The cells were pelleted, resuspended in 15 ml medium, and incubated at 37°C. At the time indicated, the cellular concentration of the analog was determined by fluorescence. Note that the efflux of analog is rapid.

FIG. 5. Monensin: Cells were incubated with 1 μ M quinacrine 17 or compound 227 for 30 min. Monensin at the indicated concentration was added either at the same time as the analog or at the end of the 30 min incubation, and the cellular concentration of the analog was determined by fluorescence.

FIG. 6. Influence of chloroquine on CpG-ODN uptake: Cells were incubated with 5 μ g/ml Texas Red CpG-ODN 1760 and indicated concentration of chloroquine 15 for the time indicated. The cells were then washed three times, and fixed. The cellular uptake of fluorescent CpG-ODN was estimated by flow cytometry. Note that chloroquine has no significant influence on CpG-ODN uptake.

FIG. 7. pH of the ODN containing compartment: WEHI 231 cells were incubated with fluorescein-labeled CpG-ODN 1760. After washing the cells were resuspended, and either 20 μ M monensin or analogs at the indicated concentration were added. 5-10 min later fluorescence was measured by flow cytometry. The change in fluorescence was converted to pH units by reference to a standard curve of fluorescence of fluorescein-labeled ODN 1760 as pH, shown on the *righthand* scale.

FIG. 8. Dimeristoylphosphatidylcholine, with number assignments used in Table 11 and FIG. 10.

FIG. 9A. NOESY spectra of compound 91 bound to phosphatidylcholine vesicles. **FIG. 9B.** Aliphatic region of the same spectra. See Table 11 for assignments, and text for details and interpretation.

FIG. 10A. Scheme 2 as described in Example 9. **FIG. 10B.** Scheme 3 as described in Example 9.

FIG. 11. Scheme 4 as described in Example 10.

FIG. 12A-E. Additional analogs with some inhibitory activity against CpG ODN immunostimulation. The level of activity is presented for each compound, and the compounds are generally presented in order of decreasing activity.

DESCRIPTION OF ILLUSTRATIVE EMBODIMENTS

The present invention is directed to therapeutic compositions and methods for inhibiting stimulation of the immune system using chloroquine, its structural analogs and derivatives, or other weak bases. These compounds and methods have application in the treatment of the following: autoimmune diseases and phenomena, transplantation rejection responses, sepsis, and other disorders affecting the immune system.

I. Treatment Uses

A. Autoimmune Diseases and Phenomena

There are numerous conditions that qualify as an autoimmune disease. They occur either when the immune system malfunctions and the lymphocytes become sensitized against self tissue cells or when self tissue cells exhibit non-self characteristics such as expression of different antigens. Some of the most common disorders are listed in Table 1, such as rheumatoid arthritis and lupus erythematosus. Other autoimmune diseases include: Alopecia Areata, Acquired Hemophilia, Ankylosing Spondylitis, Antiphospholipid Syndrome, Autoimmune Hepatitis, Autoimmune Hemolytic Anemia, Behcet's Disease, Cardiomyopathy, Celiac Sprue

Dermatitis, Chronic Fatigue Immune Dysfunction Syndrome (CFIDS), Chronic Inflammatory Demyelinating Polyneuropathy, Churg-Strauss Syndrome, Cicatricial Pemphigoid, CREST Syndrome, Cold Agglutinin Disease, Discoid Lupus, Essential Mixed Cryoglobulinemia, Fibromyalgia, Fibromyositis, Guillain-Barre, Idiopathic Pulmonary Fibrosis, Idiopathic Thrombocytopenic Purpura, IgA Nephropathy, Juvenile Arthritis, Lichen Planus, Multiple Sclerosis, Myasthenia Gravis, Polyarteritis Nodosa, Polychondritis, Polyglandular Syndromes, Dermatomyositis, Primary Agammaglobulinemia, Primary Biliary Cirrhosis, Psoriasis, Raynaud's Phenomena, Reiter's Syndrome, Sarcoidosis, Stiff-Man Syndrome, Takayasu Arthritis, Temporal Arteritis/Giant Cell Arteritis, Ulcerative Colitis, Uveitis, Vasculitis, and Vitiligo.

Table 1
Selected Autoimmune Diseases and the Targets of Their Antibodies

Systemic (Non-Organ-Specific) Diseases	Target of Antibody
Goodpasture's Syndrome	Basement membranes
Mixed Connective Tissue Disease	Nuclear ribonucleoproteins
Polymyositis	Nuclei, Jo-1, PL-7, histadyl-tRNA synthetase, threonyl-tRNA synthetase, PM-1, Mi-2
Rheumatic Fever	Myocardium, heart valves, choroid plexus
Rheumatoid Arthritis	γ -Globulin, Epstein-Barr virus-related antigens, types II and III collagen
Scleroderma	Nuclei, Scl-70, SS-A (Ro), SS-B (La), centromere
Sjögren's Syndrome	γ -Globulin, SS-A (Ro), SS-B (La)
Systemic Lupus Erythematosus	DNA, ribonucleoproteins, histones, nuclear antigens
Wegener's granulomatosis	Neutrophils
Organ-Specific Diseases	Target of Antibody
Addison's Disease	Adrenal cells
Allergic Rhinitis, Asthma, and autoimmune abnormalities	β_2 -adrenergic receptors
Autoimmune hemolytic anemia	Erythrocytes
Acquired Hemophilia	Clotting Factor VIII
Bullous Pemphigoid	Basement membrane zone of skin and mucosa

Table 1 (continued)

Systemic (Non-Organ-Specific) Diseases	Target of Antibody
Chronic Active Hepatitis	Nuclei of hepatocytes
Crohn's Disease	Lymphocytes, plasma cells, eosinophils
Glomerulonephritis	Glomeruli
Graves' Disease	Thyroid-stimulating hormone (TSH) receptor
Hashimoto's Thyroiditis	Thyroglobulin, TSH receptor
Idiopathic Hypoparathyroidism	Parathyroid cells
Idiopathic Neutropenia	Neutrophils
Idiopathic Thrombocytopenic Purpura	Platelets
Insulin-resistant Diabetes with acanthosis nigricans	Insulin receptor
Insulin-resistant Diabetes with ataxia- telangiectasia	Insulin receptor
Juvenile Insulin-dependent Diabetes	Pancreatic islet cells
Ménière's Disease	Type II collagen
Myasthenia Gravis	Acetylcholine receptors
Osteosclerosis	Type II collagen
Pemphigus	Intercellular substance of skin and mucosa
Pernicious Anemia	Gastric parietal cells, vitamin B ₁₂ binding site of intrinsic factor
Premature Ovarian Failure	Interstitial cells, corpus luteum cells
Primary Biliary Cirrhosis	Mitochondria
Spontaneous infertility	Sperm cells

5 The present invention is understood to include compounds that act to inhibit an immune response, and thus serve to allay the progression of autoimmune diseases and phenomena. These compounds comprise chloroquine and various analogs, such

as substituted versions of 4-quinolinamine, bis-4-quinolinamine, bis-9-aminoacridine, derivatives thereof, and other weak bases. Furthermore, methods of treating autoimmune diseases using these compounds are also contemplated.

5 In addition to treating autoimmune diseases, the compounds and methods of the present invention are useful generally to inhibit stimulation of the immune system. Treatment and prevention of autoimmune phenomena (caused for example by myocardial infarction), sepsis, and transplantation rejections and graft-versus-host disease are also within the scope of the present invention.

10 **B. Transplantation Rejections and Graft-Versus-Host Disease**

Transplant rejections occur as a consequence of an immune response against the transplanted organ, tissue, or cells. Antigens on the surface of the transplanted material act to signal that it is foreign, and a response ensues. Conversely, GVHD 15 occurs when the graft mounts an immune response against the host, which can happen following a bone marrow transplant. Lymphocytes in the donor marrow produce antibodies against the host and attempts to destroy cells of the host. It occurs in approximately 40% of patients receiving an allogeneic transplant.

20 Because immunostimulation occurs in both of these situations, the compounds and methods of the present invention can be used as treatments to inhibit an immune response and thus alleviate or eliminate their destructive outcomes.

C. Sepsis

25 Sepsis can be caused by many different infectious agents and microbial organisms that may or may not be involved directly with bloodstream infection. It is a condition characterized by an inflammatory response. The term "sepsis" as used herein broadly refers to conditions known as sepsis, septic shock, systemic inflammatory response syndrome (SIRS), and multiple organ dysfunction syndrome 30 (MODS). Because these conditions are caused by an inflammatory response of the immune system, the compounds and methods of the present invention can be


employed as preventative and therapeutic treatments to inhibit an immune response and reduce the incidence of sepsis.

D. Chloroquine, Chloroquine Analogs, and Other Weak Bases


5 Chloroquine and a number of its structural analogs specifically and powerfully inhibit the immunostimulatory effect of unmethylated CpG-motifs at nanomolar concentration. A large variety of analogs have been constructed, many of which possess some inhibitory effect on the immunostimulatory capacity of CpG-motifs (see FIG.1). Both chloroquine and quinacrine as well as analogs bind to duplex DNA by
10 intercalation in a relatively sequence-nonspecific fashion (Wilson, 1998). Compounds with only two fused aromatic rings (such as chloroquine) are poor intercalators and bind to DNA less avidly than quinacrine and other related tricyclic systems. The two charged groups on chloroquine and quinacrine significantly enhance their binding to DNA, and both compounds bind to duplex DNA much more
15 tightly at low salt concentrations than at higher salt.

Chloroquine, hydroxychloroquine and quinacrine also are known to induce remissions of systemic lupus erythematosus and rheumatoid arthritis by an unknown mechanism. These drugs bind to DNA by intercalation. They are weak bases and
20 partition into acidic vesicles. At high concentrations, chloroquine can collapse the pH gradient and disrupt the action of endosomal hydrolytic enzymes and the trafficking of receptors.


The compounds of the present invention include substituted 4-quinolinamines
25 (Structures A, B and C), substituted bis-4-quinolinamines (Structure D), substituted bis-9-aminoacridines (Structure E) and substituted 4-quinolinamines, substituted 9-aminoacridines combinations (Structure F).




Structure A




Structure B




Structure C



Structure D



Structure E



Structure F

In certain embodiments of the present invention, substituted 4-quinolinamines comprise substituted or unsubstituted alkyl, alkenyl or alkynyl secondary or tertiary amine groups at the 4 position (Structure A). Substituted 4-quinolinamines also may comprise a substituted or unsubstituted phenyl group (Structure B) directly attached at the 4-amine position of the quinolinamine, or can be spatially distanced from the quinolinamine 4-amine position by a lower alkyl group (Structure C). In one preferred embodiment, a hydroxy group is attached to a substituted phenyl group. Contemplated also are one or more substitutions at phenyl group positions R₉, R₁₀, R₁₁, R₁₂ and R₁₃, of Structures B and C. In preferred embodiments, the R_A group of the 4-quinolinamine is a hydrogen atom, as opposed to larger alkyl groups. The R₂ position of the substituted 4-quinolinamine compounds of the invention can be a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, an substituted or unsubstituted anthracyl group, a substituted or unsubstituted phenanthryl group or a substituted or unsubstituted styryl group. Preferably, larger aromatic groups are substituted at the R₂ position, which has been demonstrated to increase 4-quinolinamine activity.

As will be understood by those of skill in the art, small modification and changes may be made in the structure of the described substituted 4-quinolinamines, bis-4-quinolinamines, bis-9-aminoacridines and 4-quinolinamines, 9-aminoacridines combinations that inhibits immunostimulation. Furthermore, certain functional groups may be substituted for other functional groups without appreciable loss of immunosuppression. Since it is the inhibitory capacity and nature of an analog that defines its activity, certain chemical substitutions can be made in the structure and nevertheless obtain an analog with like properties. It is thus contemplated by the inventor that various changes may be made in an analog without appreciable loss of its immunosuppressive utility or activity.

II. Inhibitors of Immunostimulation

The invention involves compositions and methods that effect inhibition of immunostimulation. As used herein, the terms "inhibition of immunostimulation" or "to inhibit immunostimulation" refer to an ability to suppress or reduce, even slightly,

an immune response. An immune response can be evidenced by a number of characteristics including, but not limited to, production of lymphokines or cytokines, release of lymphokines or cytokines, proliferation of lymphocytes, activation of lymphocytes, complement fixing, induction of the complement cascade, production of antibodies, release of antibodies, release of inflammatory mediators, and binding of T cells to a T-cell receptor.

A. CpG Motifs

Bacterial DNA is increasingly recognized as a powerful modulator of immunity (Krieg, 1996, *Trends in Microbiol.*, 4:73-6), stimulating the polyclonal proliferation of B-cells and the production of cytokines by monocytes and other cells (Ballas *et al.*, 1996, *J. Immunol.*, 157:1840-5). This activity is attributed to unmethylated CpG sequences in bacterial DNA, which are uncommon in vertebrate DNA (Krieg *et al.*, 1995, *Nature*, 374:546-9). Single stranded oligodeoxynucleotides that have the motif PuPuCGPyPy (CpG-ODN) mimic many of the actions of bacterial DNA (Krieg *et al.*, 1996, *Antisense & Nucleic Acid Drug Devel.*, 6:133-9), and powerfully inhibit the induction of apoptosis in mouse WEHI 231 B-cells by anti-surface IgM (Yi *et al.*, 1996, *J. Immunol.*, 157:4918-25; Macfarlane *et al.*, 1997, *Immunology* 91:386). This system is a convenient and reproducible model to study responses to immunostimulatory DNA.

The term "CpG" refers to a nucleic acid sequence or molecule having a cytosine connected by a phosphorous-containing link to the 5' end of a guanosine. The term "methylated CpG" refers to the methylation of the cytosine on the pyrimidine ring, usually occurring at the 5-position of the pyrimidine ring. The term "CpG-motif" refers to a nucleic acid sequence or molecule having CpG in a context that is immunostimulatory, especially when the CpG is adjacent to two purines on its 5' end and two pyrimidines on its 3' end.

Bacterial DNA immobilized on beads does not stimulate immune responses, suggesting that internalization of the DNA is required for activity. DNA and

oligodeoxynucleotides are endocytosed into acidic vesicles, and are then transported to the cytoplasm and nucleus of cells.

The mechanism of inhibition by chloroquine and its analogs was examined using 4-aminoquinolines, quinacrine, 9-aminoacridines, and novel dibasic analogs, many of which are fluorescent. WEHI 231 murine B-lymphoma cells accumulated analogs up to a concentration several hundred-fold higher than the medium. Uptake was rapid, nonsaturable, reversible, and partially inhibited by monensin, an agent that collapses pH gradients within cells. Uptake did not correlate highly with efficacy as inhibitors of CpG-ODN-induced effects, suggesting that analogs act by a specific action. Confocal microscopy revealed analogs concentrating in large peripheral organelles. CpG-ODN is taken up into acidified, small, perinuclear vesicles by cells, which uptake is believed to be necessary for immunostimulatory activity. This cellular uptake of fluorescent CpG-ODN is not inhibited by the analogs. The pH of intracellular CpG-ODN (6.4) was not affected by analogs at the concentration required for inhibition, but pH was increased by higher concentrations. UV-spectroscopy revealed no binding of analogs to CpG-ODN. Nuclear Overhauser effect spectroscopy revealed that an analog bound to phosphatidylcholine vesicles, with the ring structure of the analog buried within the lipid and the side-chain facing the aqueous environment. The conclusion is that the analogs do not inhibit the action of CpG-ODN by preventing the uptake or acidification of CpG-ODN.

B. Linkers/Coupling Agents

If desired, dimers or multimers of the chloroquine analogs may be joined *via* a biologically-releasable bond, such as a selectively-cleavable linker or amino acid sequence. For example, peptide linkers that include a cleavage site for an enzyme preferentially located or active within a tumor environment are contemplated. Exemplary forms of such peptide linkers are those that are cleaved by urokinase, plasmin, thrombin, Factor IXa, Factor Xa, or a metalloproteinase, such as collagenase, gelatinase, or stromelysin.

Additionally, while numerous types of disulfide-bond containing linkers are known which can successfully be employed to conjugate the toxin moiety with the targeting agent, certain linkers will generally be preferred over other linkers, based on differing pharmacologic characteristics and capabilities. For example, linkers that contain a disulfide bond that is sterically "hindered" are to be preferred, due to their greater stability *in vivo*, thus preventing release of the toxin moiety prior to binding at the site of action.

Cross-linking reagents are used to form molecular bridges that tie together functional groups of two different molecules, e.g., a stabilizing and coagulating agent. However, it is contemplated that dimers or multimers of the same analog can be made or that heteromeric complexes comprised of different analogs can be created. To link two different compounds in a step-wise manner, hetero-bifunctional cross-linkers can be used that eliminate unwanted homopolymer formation.

TABLE 2
HETERO-BIFUNCTIONAL CROSS-LINKERS

linker	Reactive Toward	Advantages and Applications	Spacer Arm Length\after cross-linking
SMPT	Primary amines Sulphydryls	· Greater stability	11.2 Å
SPDP	Primary amines Sulphydryls	· Thiolation · Cleavable cross-linking	6.8 Å
LC-SPDP	Primary amines Sulphydryls	· Extended spacer arm	15.6 Å
Sulfo-LC-SPDP	Primary amines Sulphydryls	· Extended spacer arm · Water-soluble	15.6 Å
SMCC	Primary amines Sulphydryls	· Stable maleimide reactive group · Enzyme-antibody conjugation · Hapten-carrier protein conjugation	11.6 Å
Sulfo-SMCC	Primary amines Sulphydryls	· Stable maleimide reactive group · Water-soluble · Enzyme-antibody conjugation	11.6 Å
MBS	Primary amines Sulphydryls	· Enzyme-antibody conjugation · Hapten-carrier protein conjugation	9.9 Å
Sulfo-MBS	Primary amines Sulphydryls	· Water-soluble	9.9 Å
SIAB	Primary amines Sulphydryls	· Enzyme-antibody conjugation	10.6 Å
Sulfo-SIAB	Primary amines Sulphydryls	· Water-soluble	10.6 Å
SMPB	Primary amines Sulphydryls	· Extended spacer arm · Enzyme-antibody conjugation	14.5 Å
Sulfo-SMPB	Primary amines Sulphydryls	· Extended spacer arm · Water-soluble	14.5 Å
EDC/Sulfo-NHS	Primary amines Carboxyl groups	· Hapten-Carrier conjugation	0
ABH	Carbohydrates Nonselective	· Reacts with sugar groups	11.9 Å

An exemplary hetero-bifunctional cross-linker contains two reactive groups: one reacting with primary amine group (*e.g.*, N-hydroxy succinimide) and the other reacting with a thiol group (*e.g.*, pyridyl disulfide, maleimides, halogens, etc.). Through the primary amine reactive group, the cross-linker may react with the lysine residue(s) of one protein (*e.g.*, the selected antibody or fragment) and through the thiol reactive group, the cross-linker, already tied up to the first protein, reacts with the cysteine residue (free sulfhydryl group) of the other protein (*e.g.*, the selective agent).

It is preferred that a cross-linker having reasonable stability in blood will be employed. Numerous types of disulfide-bond containing linkers are known that can be successfully employed to conjugate targeting and therapeutic/preventative agents. Linkers that contain a disulfide bond that is sterically hindered may prove to give greater stability *in vivo*, preventing release of the targeting peptide prior to reaching the site of action. These linkers are thus one group of linking agents.

Another cross-linking reagents for use in immunotoxins is SMPT, which is a bifunctional cross-linker containing a disulfide bond that is "sterically hindered" by an adjacent benzene ring and methyl groups. It is believed that steric hindrance of the disulfide bond serves a function of protecting the bond from attack by thiolate anions such as glutathione which can be present in tissues and blood, and thereby help in preventing decoupling of the conjugate prior to the delivery of the attached agent to the target site.

The SMPT cross-linking reagent, as with many other known cross-linking reagents, lends the ability to cross-link functional groups such as the SH of cysteine or primary amines (*e.g.*, the epsilon amino group of lysine). Another possible type of cross-linker includes the hetero-bifunctional photoreactive phenylazides containing a cleavable disulfide bond such as sulfosuccinimidyl-2-(p-azido salicylamido) ethyl-1,3'-dithiopropionate. The N-hydroxy-succinimidyl group reacts with primary amino groups and the phenylazide (upon photolysis) reacts non-selectively with any amino acid residue.

In addition to hindered cross-linkers, non-hindered linkers also can be employed in accordance herewith. Other useful cross-linkers, not considered to contain or generate a protected disulfide, include SATA, SPDP and 2-iminothiolane (Wawrzynczak & Thorpe, 1987). The use of such cross-linkers is well understood in the art.

In preferred embodiments utilizing linkers, flexible linkers are employed to create conjugates.

Once conjugated, the analog generally will be purified to separate the conjugate from unconjugated analog or coagulants and from other contaminants. A large number of purification techniques are available for use in providing conjugates of a sufficient degree of purity to render them clinically useful. Purification methods based upon size separation, such as gel filtration, gel permeation or high performance liquid chromatography, will generally be of most use.

C. Combined Therapy with Analogs and Traditional Treatment

In many therapies, it will be advantageous to provide more than one functional therapeutic. Such "combined" therapies may have particular import in treating aspects of autoimmune diseases/phenomena and tissue/organ rejections. Thus, one aspect of the present invention utilizes at least one chloroquine analog compound for treatment of immunostimulation, while a second therapy also is provided.

The non-targeted treatment may precede or follow the targeted agent treatment by intervals ranging from minutes to weeks. In embodiments where the other agent and analogs are applied separately to the cell, one would generally ensure that a significant period of time did not expire between the time of each delivery, such that the agent and analog would still be able to exert an advantageously combined effect on the cell. In such instances, it is contemplated that one would contact the cell with both modalities within about 12-24 hours of each other and, more preferably, within about 6-12 hours of each other, with a delay time of only about 12 hours being most preferred. In some

situations, it may be desirable to extend the time period for treatment significantly, however, where several days (2, 3, 4, 5, 6 or 7) to several weeks (1, 2, 3, 4, 5, 6, 7 or 8) lapse between the respective administrations.

- 5 It also is conceivable that more than one administration of either agent will be desired. Various combinations may be employed, where the analog is "A" and the non-analog is "B", as exemplified below:

A/B/A B/A/B B/B/A A/A/B B/A/A A/B/B B/B/B/A B/B/A/B

10

A/A/B/B A/B/A/B A/B/B/A B/B/A/A B/A/B/A B/A/A/B B/B/B/A

A/A/A/B B/A/A/A A/B/A/A A/A/B/A A/B/B/B B/A/B/B B/B/A/B

- 15 Other combinations are contemplated. For example, in the context of the present invention, it is contemplated that the present invention's compounds and methods of inhibiting immunostimulation could be used in conjunction with non-analog agents, including steroidal treatment. To inhibit immunostimulation using the methods and compositions of the present invention, one would generally contact a "target" cell with
- 20 at least one analog and at least one other agent; these compositions would be provided in a combined amount effective achieve these goals. This process may involve contacting the cells with an analog and the agent(s) or factor(s) at the same time. This may be achieved by contacting the cell with a single composition or pharmacological formulation that includes both agents, or by contacting the cell with two distinct
- 25 compositions or formulations, at the same time, wherein one composition includes an analog and the other includes the agent.

- Agents or factors suitable for use in a combined therapy are any chemical compound or treatment method with immunosuppressive activity; therefore, the term
- 30 "immunosuppressive agent" that is used throughout this application refers to an agent with immunosuppressive activity. Immunosuppressive agents such as azathioprine and cyclosporin are employed in transplant procedures to treat and prevent rejections.

Compounds or methods used to treat GVHD include corticosteroids such as prednisone, antithymocyte globulins, cyclosporine A, cyclophosphamide, and methotrexate. Thalidomide is occasionally employed in combination with one of the previously mentioned corticosteroids to treat GVHD, and it is contemplated that thalidomide could also be used in combination with the compounds of the present invention. Similarly, patients with autoimmune diseases are administered immunosuppressant medications like corticosteroids, cyclophosphamide, and azathioprine.

In the treatment of sepsis, other agents or compounds that would be useful for use in a combination therapy with the compounds and methods of the claimed invention include antibiotics such as cephalosporin, fluoroquinolones, penicillins, carbapenems, β -lactams- β -lactamase inhibitors, ampicillin, vancomycin, metronidazole, clindamycin, and trovafloxacin, as well as with corticosteroids, vasopressor agents, vasoconstrictors, and beta agonists.

The skilled artisan is directed to "Remington's Pharmaceutical Sciences" 15th Edition, chapter 33, in particular pages 624-652. Some variation in dosage will necessarily occur depending on the condition of the subject being treated. The person responsible for administration will, in any event, determine the appropriate dose for the individual subject. Moreover, for human administration, preparations should meet sterility, pyrogenicity, and general safety and purity standards as required by FDA Office of Biologics standards.

The inventors propose that local, regional delivery of at least one analog will be a very efficient method for delivering a therapeutically effective compound to counteract the clinical disease. Similarly, the immunosuppressive agent may be directed to a particular, affected region of the subject's body. Alternatively, systemic delivery of compounds and/or the agents may be appropriate in certain circumstances, for example, where extensive tissue damage has occurred.

III. Therapeutic Formulations and Routes of Administration

The present invention discloses the compositions and methods involving a CpG analog that inhibits stimulation of the immune system. Where clinical applications are contemplated, it will be necessary to prepare the compositions of the present invention as pharmaceutical compositions, *i.e.*, in a form appropriate for *in vivo* applications. Generally, this will entail preparing compositions that are essentially free of pyrogens, as well as other impurities that could be harmful to humans.

A. Formulations

The compounds of the present invention may be operatively linked or attached to a selective agent or compound. Different and varied therapeutic compounds are illustrated, which include chloroquines and its analogs and derivatives. Liposomes and carrier molecules containing any of the foregoing are also contemplated.

One will generally desire to employ appropriate salts and buffers to render compositions stable and allow for uptake by target cells. Buffers also will be employed when recombinant cells are introduced into a patient. Aqueous compositions of the present invention comprise an effective amount of the vector to cells, dissolved or dispersed in a pharmaceutically acceptable carrier or aqueous medium. Such compositions also are referred to as inocula. The phrase "pharmaceutically or pharmacologically acceptable" refer to molecular entities and compositions that do not produce adverse, allergic, or other untoward reactions when administered to an animal or a human. As used herein, "pharmaceutically acceptable carrier" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents and the like. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the vectors or cells of the present invention, its use in therapeutic compositions is contemplated. Supplementary active ingredients also can be incorporated into the compositions.

The active compositions of the present invention include classic pharmaceutical preparations. Administration of these compositions according to the present invention will be *via* any common route so long as the target tissue is available *via* that route. This includes oral, nasal, buccal, rectal, vaginal or topical. Alternatively, administration may be by orthotopic, intradermal, subcutaneous, intramuscular, intraperitoneal or intravenous injection. Such compositions would normally be administered as pharmaceutically acceptable compositions, described *supra*.

The active compounds may be administered *via* any suitable route, including parenterally or by injection or inhalation. Solutions of the active compounds as free base or pharmacologically acceptable salts can be prepared in water suitably mixed with a surfactant, such as hydroxypropylcellulose. Dispersions also can be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases the form must be sterile and must be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms, such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), suitable mixtures thereof, and vegetable oils. The proper fluidity can be maintained, for example, by the use of a coating, such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars or sodium chloride. Prolonged absorption of the

injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

5 Sterile injectable solutions are prepared by incorporating the active compounds in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the various sterilized active ingredients into a sterile vehicle which contains the basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the
10 preparation of sterile injectable solutions, the preferred methods of preparation are vacuum-drying and freeze-drying techniques which yield a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

15 As used herein, "pharmaceutically acceptable carrier" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents and the like. The use of such media and agents for pharmaceutical active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active ingredient, its use in the
20 therapeutic compositions is contemplated. Supplementary active ingredients also can be incorporated into the compositions.

The compositions of the present invention may be formulated in a neutral or salt form. Pharmaceutically-acceptable salts include the acid addition salts (formed
25 with the free amino groups of the protein) and which are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric, mandelic, and the like. Salts formed with the free carboxyl groups also can be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as
30 isopropylamine, trimethylamine, histidine, procaine and the like.

Compositions may be conventionally administered parenterally, by injection, for example, either subcutaneously or intramuscularly. Additional formulations which are suitable for other modes of administration include suppositories and, in some cases, oral formulations. For suppositories, traditional binders and carriers may include, for example, polyalkylene glycols or triglycerides: such suppositories may be formed from mixtures containing the active ingredient in the range of about 0.5% to about 10%, preferably about 1 to about 2%. Oral formulations include such normally employed excipients as, for example, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate and the like. These compositions take the form of solutions, suspensions, tablets, pills, capsules, sustained release formulations or powders and contain about 10 to about 95% of active ingredient, preferably about 25 to about 70%.

1. *Liposomes as Carriers of Selected Compounds*

In one embodiment of the invention, the selected compound of the present invention may be entrapped in a liposome. Liposomes are vesicular structures characterized by a phospholipid bilayer membrane and an inner aqueous medium. Multilamellar liposomes have multiple lipid layers separated by aqueous medium. They form spontaneously when phospholipids are suspended in an excess of aqueous solution. The lipid components undergo self-rearrangement before the formation of closed structures and entrap water and dissolved solutes between the lipid bilayers (Ghosh and Bachhawat, 1991). Also contemplated are cationic lipid-nucleic acid complexes, such as lipofectamine-nucleic acid complexes.

"Liposome" is a generic term encompassing a variety of single and multilamellar lipid vehicles formed by the generation of enclosed lipid bilayers. Phospholipids are used for preparing the liposomes according to the present invention and can carry a net positive charge, a net negative charge or are neutral. Dicyetyl phosphate can be employed to confer a negative charge on the liposomes, and stearylamine can be used to confer a positive charge on the liposomes.

Lipids suitable for use according to the present invention can be obtained from commercial sources. For example, dimyristyl phosphatidylcholine ("DMPC") can be obtained from Sigma Chemical Co., dicetyl phosphate ("DCP") is obtained from K & K Laboratories (Plainview, NY); cholesterol ("Chol") is obtained from Calbiochem-Behring; dimyristyl phosphatidylglycerol ("DMPG") and other lipids may be obtained from Avanti Polar Lipids, Inc. (Birmingham, Ala.). Stock solutions of lipids in chloroform, chloroform/methanol or *t*-butanol can be stored at about -20°C. Preferably, chloroform is used as the only solvent since it is more readily evaporated than methanol.

Phospholipids from natural sources, such as egg or soybean phosphatidylcholine, brain phosphatidic acid, brain or plant phosphatidylinositol, heart cardiolipin and plant or bacterial phosphatidylethanolamine are preferably not used as the primary phosphatide, *i.e.*, constituting 50% or more of the total phosphatide composition, because of the instability and leakiness of the resulting liposomes.

Liposomes used according to the present invention can be made by different methods. The size of the liposomes varies depending on the method of synthesis. A liposome suspended in an aqueous solution is generally in the shape of a spherical vesicle, having one or more concentric layers of lipid bilayer molecules. Each layer consists of a parallel array of molecules represented by the formula XY, wherein X is a hydrophilic moiety and Y is a hydrophobic moiety. In aqueous suspension, the concentric layers are arranged such that the hydrophilic moieties tend to remain in contact with an aqueous phase and the hydrophobic regions tend to self-associate. For example, when aqueous phases are present both within and without the liposome, the lipid molecules will form a bilayer, known as a lamella, of the arrangement XY-YX.

Liposomes within the scope of the present invention can be prepared in accordance with known laboratory techniques. In one preferred embodiment, liposomes are prepared by mixing liposomal lipids, in a solvent in a container, *e.g.*, a glass, pear-shaped flask. The container should have a volume ten-times greater than the volume of the expected suspension of liposomes. Using a rotary evaporator, the solvent

is removed at approximately 40°C under negative pressure. The solvent normally is removed within about 5 min to 2 hours, depending on the desired volume of the liposomes. The composition can be dried further in a desiccator under vacuum. The dried lipids generally are discarded after about 1 week because of a tendency to deteriorate with time.

Dried lipids can be hydrated at approximately 25-50 mM phospholipid in sterile, pyrogen-free water by shaking until all the lipid film is resuspended. The aqueous liposomes can be then separated into aliquots, each placed in a vial, lyophilized and sealed under vacuum.

In the alternative, liposomes can be prepared in accordance with other known laboratory procedures: the method of Bangham *et al.* (1965), the contents of which are incorporated herein by reference; the method of Gregoriadis, as described in *DRUG CARRIERS IN BIOLOGY AND MEDICINE*, G. Gregoriadis ed. (1979) pp. 287-341, the contents of which are incorporated herein by reference; the method of Deamer and Uster (1983), the contents of which are incorporated by reference; and the reverse-phase evaporation method as described by Szoka and Papahadjopoulos (1978). The aforementioned methods differ in their respective abilities to entrap aqueous material and their respective aqueous space-to-lipid ratios.

The dried lipids or lyophilized liposomes prepared as described above may be reconstituted in a solution of nucleic acid and diluted to an appropriate concentration with an suitable solvent, *e.g.*, DPBS. The mixture is then vigorously shaken in a vortex mixer. Unencapsulated nucleic acid is removed by centrifugation at $29,000 \times g$ and the liposomal pellets washed. The washed liposomes are resuspended at an appropriate total phospholipid concentration, *e.g.*, about 50-200 mM. The amount of nucleic acid encapsulated can be determined in accordance with standard methods. After determination of the amount of nucleic acid encapsulated in the liposome preparation, the liposomes may be diluted to appropriate concentration and stored at 4°C until use.

In another embodiment, the lipid dioleoylphosphatidylcholine is employed. Nuclease-resistant oligonucleotides were mixed with lipids in the presence of excess t-butanol. The mixture was vortexed before being frozen in an acetone/dry ice bath. The frozen mixture was lyophilized and hydrated with Hepes-buffered saline (1 mM
5 Hepes, 10 mM NaCl, pH 7.5) overnight, and then the liposomes were sonicated in a bath type sonicator for 10 to 15 min. The size of the liposomal-oligonucleotides typically ranged between 200-300 nm in diameter as determined by the submicron particle sizer autodilute model 370 (Nicomp, Santa Barbara, CA).

10 Liposomes are formed from phospholipids that are dispersed in an aqueous medium and spontaneously form multilamellar concentric bilayer vesicles. Liposomes bear many resemblances to cellular membranes and are contemplated for use in connection with the present invention as drug delivery agents.

15 The formation and use of liposomes is generally known to those of skill in the art. For example, several U.S. Patents concern the preparation and use of liposomes that encapsulate biologically active materials, *e.g.*, U.S. Patent 4,485,054; 4,089,801; 4,234,871; and 4,016,100; each incorporated herein by reference. Mostly, it is contemplated that intravenous injection of liposomal preparations would be used, but
20 pessaries are also possible.

B. Routes of Administration

Upon formulation, solutions will be administered in a manner compatible with the dosage formulation and in such amount as is therapeutically effective. The
25 formulations are easily administered in a variety of dosage forms such as injectable solutions, drug release capsules and the like. For parenteral administration in an aqueous solution, for example, the solution should be suitably buffered if necessary and the liquid diluent first rendered isotonic with sufficient saline or glucose. These particular aqueous solutions are especially suitable for intravenous, intramuscular,
30 subcutaneous and intraperitoneal administration.

1. *Nasal Administration*

One may also use nasal solutions or sprays, aerosols or inhalants in the present invention. Nasal solutions are usually aqueous solutions designed to be administered to the nasal passages in drops or sprays. Nasal solutions are prepared so that they are
5 similar in many respects to nasal secretions, so that normal ciliary action is maintained. Thus, the aqueous nasal solutions usually are isotonic and slightly buffered to maintain a pH of 5.5 to 6.5.

In addition, antimicrobial preservatives, similar to those used in ophthalmic
10 preparations, and appropriate drug stabilizers, if required, may be included in the formulation. Various commercial nasal preparations are known and include, for example, antibiotics and antihistamines and are used for asthma prophylaxis.

2. *Oral Administration*

15 In certain embodiments, active compounds may be administered orally. This is contemplated to be useful as many substances contained in tablets designed for oral use are absorbed by mucosal epithelia along the gastrointestinal tract.

Also, if desired, the peptides, antibodies and other agents may be rendered
20 resistant, or partially resistant, to proteolysis by digestive enzymes. Such compounds are contemplated to include chemically designed or modified agents; dextrorotatory peptides; and peptide and liposomal formulations in time release capsules to avoid peptidase and lipase degradation.

25 For oral administration, the active compounds may be administered, for example, with an inert diluent or with an assimilable edible carrier, or they may be enclosed in hard or soft shell gelatin capsule, or compressed into tablets, or incorporated directly with the food of the diet. For oral therapeutic administration, the active compounds may be incorporated with excipients and used in the form of ingestible
30 tablets, buccal tables, troches, capsules, elixirs, suspensions, syrups, wafers, and the like.

The oral compositions and preparations should contain at least 0.1% of active compound. The percentage of the compositions and preparations may, of course, be varied and may conveniently be between about 2 to about 60% of the weight of the unit. The amount of active compounds in such therapeutically useful compositions is such
5 that a suitable dosage will be obtained.

The tablets, troches, pills, capsules and the like may also contain the following: a binder, as gum tragacanth, acacia, cornstarch, or gelatin; excipients, such as dicalcium phosphate; a disintegrating agent, such as corn starch, potato starch, alginic acid and the
10 like; a lubricant, such as magnesium stearate; and a sweetening agent, such as sucrose, lactose or saccharin may be added or a flavoring agent, such as peppermint, oil of wintergreen, or cherry flavoring. When the dosage unit form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier.

15 Various other materials may be present as coatings or to otherwise modify the physical form of the dosage unit. For instance, tablets, pills, or capsules may be coated with shellac, sugar or both. A syrup or elixir may contain the active compounds sucrose as a sweetening agent methyl and propylparabens as preservatives, a dye and flavoring, such as cherry or orange flavor. Of course, any material used in preparing any dosage
20 unit form should be pharmaceutically pure and substantially non-toxic in the amounts employed. In addition, the active compounds may be incorporated into sustained-release preparation and formulations.

Upon formulation, the compounds will be administered in a manner compatible
25 with the dosage formulation and in such amount as is therapeutically effective. The formulations are easily administered in a variety of dosage forms, as described herein.

3. *Pessaries*

Additional formulations which are suitable for other modes of administration
30 include vaginal suppositories and pessaries. A rectal pessary or suppository may also be used.

Suppositories are solid dosage forms of various weights and shapes, usually medicated, for insertion into the rectum, vagina or the urethra. After insertion, suppositories soften, melt or dissolve in the cavity fluids.

5 In general, for suppositories, traditional binders and carriers may include, for example, polyalkylene glycols or triglycerides; such suppositories may be formed from mixtures containing the active ingredient in the range of 0.5% to 10%, preferably 1%-2%.

10 Vaginal suppositories or pessaries are usually globular or oviform and weighing about 5 g each. Vaginal medications are available in a variety of physical forms, *e.g.*, creams, gels or liquids, which depart from the classical concept of suppositories. Vaginal tablets, however, do meet the definition, and represent convenience both of administration and manufacture.

15 "Unit dose" is defined as a discrete amount of a therapeutic composition dispersed in a suitable carrier. In this connection, sterile aqueous media which can be employed will be known to those of skill in the art in light of the present disclosure. For example, a unit dose could be dissolved in 1 ml of isotonic NaCl solution and
20 either added to 1000 ml of hypodermoclysis fluid or injected at the proposed site of infusion, (see for example, "Remington's Pharmaceutical Sciences" 15th Edition, pages 1035-1038 and 1570-1580). Some variation in dosage will necessarily occur depending on the condition of the subject being treated. The person responsible for administration will, in any event, determine the appropriate dose for the individual
25 subject. Moreover, for human administration, preparations should meet sterility, pyrogenicity, general safety and purity standards as required by FDA Office of Biologics standards.

B. Kits

All the essential materials and reagents required for delivering analogs and agents to effect inhibition of immunostimulation may be assembled together in a kit. This generally will comprise selected analogs prepared in an adminsterable formulation
5 or prepared for use as an adminsterable formulation. Such kits will comprise distinct containers for each individual reagent.

When the components of the kit are provided in one or more liquid solutions, the liquid solution preferably is an aqueous solution, with a sterile aqueous solution being
10 particularly preferred. For *in vivo* use, the expression construct may be formulated into a pharmaceutically acceptable syringeable composition. In this case, the container means may itself be an inhalent, syringe, pipette, eye dropper, or other such like apparatus, from which the formulation may be applied to an infected area of the body, such as the lungs, injected into a subject, or even applied to and mixed with the other
15 components of the kit.

The components of the kit may also be provided in dried or lyophilized forms. When reagents or components are provided as a dried form, reconstitution generally is by the addition of a suitable solvent. It is envisioned that the solvent also may be
20 provided in another container means.

The kits of the present invention also will typically include a means for containing the vials in close confinement for commercial sale such as, *e.g.*, injection or blow-molded plastic containers into which the desired vials are retained.
25

Irrespective of the number or type of containers, the kits of the invention also may comprise, or be packaged with, an instrument for assisting with the injection/administration or placement of the ultimate complex composition within the body of an animal. Such an instrument may be an inhalant, syringe, pipette, forceps,
30 measured spoon, eye dropper or any such medically approved delivery vehicle.

IV. Examples

The following examples are included to demonstrate preferred embodiments of the invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples which follow represent techniques discovered by the inventors to function well in the practice of the invention, and thus can be considered to constitute preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments which are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.


EXAMPLE 1: Structure-Activity Relationship Analysis of Substituted 4-Amioquinolinamines

A detailed structure-activity relationship (SAR) analysis of quinoline antagonists of immunostimulatory CpG-ODNs was undertaken. The structure of compounds 1-32 are shown in Tables 4-9. The synthesis work together with SAR analysis of the synthesized quinolines culminated in the finding of an extremely active agent 32.

Substituted quinolines 1 (Strekowski *et al.*, 1996), 3 (Strekowski *et al.*, 1996), 4 (Strekowski *et al.*, 1991), 6 (Strekowski *et al.*, 1996) (Table 4), 7 (Strekowski *et al.*, 1996), 9 (Strekowski *et al.*, 1996), 10 (Strekowski *et al.*, 1996) (Table 5), and 25 (Strekowski *et al.*, 1997) (Table 7) were available from other studies. Compounds 2 and 5 were obtained by using a general procedure (Strekowski *et al.*, 1992). Compounds 11 and 14 (Table 6) were obtained by treatment of 4-chloroquinoline with the corresponding amine (Strekowski *et al.*, 1997). An efficient synthesis of 2-aryl-4-chloroquinolines has recently been reported (Strekowski *et al.*, 1997). These compounds served as substrates for 8, 12, 13, 15, and 16, 20-24, and 26-32 (Table 8). This chemistry is illustrated in Scheme 1 for the preparation of 28 by the reaction of 4-chloro-2-naphthylquinoline with 4-hydroxyaniline. A subsequent Mannich reaction of 28 with formaldehyde and N-methylpiperazine furnished 32. Compounds 29-31 were prepared in a similar manner. Finally, the quaternary compounds 17-19 (Table 6) were synthesized by the reaction of the corresponding aminoquinolines with MeI in

DMF. The initially formed iodide of **17** [**15** (0.12 mmol), MeI (0.10 mmol), Na₂CO₃ (0.24 mmol), DMF (2 mL), stirring at 23 °C for 12 h] resisted all attempts of crystallization. A bromide/ hydrobromide derivative of **17** crystallized following treatment of a solution of the iodide in MeOH with HBr (48%, 3 mmol). Dibromides **18** and **19** were prepared in a similar way by the reaction of the respective quinolines **15** and **14** (0.12 mmol) with MeI (2 mmol) followed by treatment with HBr. Solid quinolines were purified by crystallization, and oily derivatives were transformed into solid salts by using a general procedure (Strekowski *et al.*, 1991) and the salts were crystallized. All compounds gave satisfactory results of elemental analysis and their structures were fully consistent with ¹H and ¹³C NMR data. (Compound, mp (crystallization solvent): **2**²HBr²H₂O, 264-267 °C (95% EtOH); **5**²HBr, 231-233 °C (95% EtOH); **8**, 52-54 °C (CH₂Cl₂/Et₂O); **11**²H₃PO₄²H₂O, 222-225 °C (95% MeOH); **12**²HBr²H₂O, 243-245 °C (95% MeOH); **13**²HBr²H₂O, 249-250 °C (95% MeOH); **14**²HCl²H₂O, 144-147 °C (95% EtOH); **15**²HBr, 205-207 °C (95% MeOH); **17**, 214-216 °C (30% MeOH); **18**, 209-212 (acetone); **19**²H₂O, 239-241 °C (95% EtOH); **20**, 87-89 °C (AcOEt); **21**, 65-67 °C (AcOEt); **22**, 189-190 °C (Et₂O); **23**²HBr, 220-222 °C (95% MeOH); **24**²HCl²H₂O, 122-129 °C (95% MeOH); **26**²H₂O, 173-175 °C (95% MeOH); **27**²HCl, 267-272 °C (95% MeOH); **28**²HCl^{0.5}H₂O, >300 °C (95% EtOH); **29**³HCl^{2.5}H₂O, 224-226 °C (95% EtOH); **30**⁴H₂O, 161-168 °C (95% MeOH); **31**³HCl²H₂O, 213-216 °C (95% EtOH/*t*-BuOMe); **32**⁵HCl⁵H₂O, 218-222 °C (95% EtOH). ¹H NMR for **32**⁵HCl⁵H₂O (DMSO-*d*₆, 400 MHz): δ 2.74 (s, 6H), 3.35-3.75 (m, 16H), 4.45 (s, 4H), 7.17 (s, 1H), 7.67 (m, 2H), 7.81 (t, J = 8 Hz, 1H), 7.87 (s, 2H), 8.02-8.23 (m, 5H), 8.46 (d, J = 8 Hz, 1H), 8.82 (s, 1H), 8.95 (d, J = 8 Hz, 1H), 11.21 (s, exchangeable with D₂O, 1H), 12.05 (br s, exchangeable with D₂O, 1H). ¹³C NMR for **32**⁵HCl⁵H₂O (DMSO-*d*₆, 75 MHz): δ 41.8, 47.5, 49.9, 54.0, 99.4, 116.3, 119.7, 120.6, 123.5, 125.1, 127.2, 127.4, 127.8, 128.5, 128.8, 129.0, 129.2, 129.4, 129.5, 131.2, 132.4, 134.2, 134.3, 138.9, 152.9, 154.7, 155.4. Spectral data for other compounds will be published in due course).

Scheme 1




- (i): $p\text{-H}_2\text{NC}_6\text{H}_4\text{OH}$ (3 equiv), 130 °C, 4 h, then silica gel chromatography (Et_2O); (ii): **28** (1 mmol), *N*-methylpiperazine (5 mmol), aq CH_2O (13 M, 5 mmol),
5 EtOH (20 mL), reflux, 24 h; then silica gel chromatography ($\text{AcOEt}/\text{Et}_3\text{N}$, 3/2).

The SAR studies were initiated following the finding that a 2-(2-naphthyl)quinoline **1** is more active than chloroquine and 2-phenylchloroquine (Table 4). It was of interest, therefore, to analyze analogs of **1** containing different groups at
10 position 2 of the quinoline. In comparison to **1**, the activity remains unchanged for **2**, which contains a larger 3-phenanthryl group and is slightly decreased for the 1-naphthyl derivative **3** with a severe steric hindrance around the inter-ring bond. Interesting results are the high efficacy of quinoline **4** that is substituted with a relatively small *para*-tolyl group and an even greater activity for compound **5**, which
15 contains a styryl moiety. Comparison of compounds **4** and **6** reveals that the electron-withdrawing trifluoromethyl group in **6** exerts a strongly negative effect on the activity.


Table 4. Activities of Chloroquine,^a 2-Phenylchloroquine,^a and 2-Substituted 4-[2-(Dimethylamino)ethyl]quinolines 1-6

No	Ar	EC ₅₀ (nM)
1	2-naphthyl	9.1
2	3-phenanthryl	11.0
3	1-naphthyl	39.8
4	4-MePh	12.3
5	<i>trans</i> -CH=CHPh	6.5
6	4-CF ₃ Ph	155



Chloroquine: R=H (EC₅₀ = 110 nM)^a
 2-Ph-chloroquine: R = Ph (EC₅₀ = 51.3)


^aTaken from MacFarlane *et al.*, 1998. Compounds 1-6 were assayed under the same conditions. The EC₅₀ is the concentration required for half-maximal inhibition of CpG-ODN effect on thymidine uptake by WEHI 231 B-cells in the presence of α-sIgM. The estimated experimental error is ±15%.



1-6

The effect of basicity of the ring nitrogen atom in selected quinolines **1** and **7–10** was investigated (Table 5). As discussed previously (Strekowski *et al.*, 1996), the given pK_a values are functions of electronic effects of the 4-substituent at the quinoline including inhibition of the conjugation effect in **7** due to steric hindrance. As can be seen from Table 5, the pK_a values of **1**, **7–10** parallel nicely the respective EC_{50} values.

Table 5. The Experimental pK_a Values of Quinolines **1**, **7–10**^a and Their Activity



No.	R	pK_a	EC_{50}
(nM)			
1	$NH(CH_2)_2NMe_2$	7.1	9.1
7	$N(Me)(CH_2)_2NMe_2$	6.2	416
8	$O(CH_2)_2NMe_2$	6.1	478
9	$S(CH_2)_2NMe_2$	4.4	2400
10	$C(O)NH(CH_2)_2NMe_2$	2.9	>10,000


^aThe values for **1**, **7**, **9**, and **10** are taken from Strekowski *et al.*, *J. Med. Chem.* 1996.

Further analogs of **1** are analyzed in Table 6. These are quinolines with and without aryl substitution, with an increasing length of the (dimethylamino)polymethylene chain, and several quaternized derivatives in which a positive charge is permanently fixed. There is a dramatic increase in activity, due to 2-aryl substitution, as can be seen from comparison of **11** to **12** and **13** and comparison of **14** to **15**. On the other hand, the activities of compounds **4** and **1** containing the same (dimethylamino)dimethylene side chain are similar to those of the respective (dimethylamino)trimethylene analogs **12** and **13**. As the chain length

increases in the series of compounds **13**, **15**, and **16**, which contain the same 2-(2-naphthyl)quinolin-4-amine core, the activity reaches a maximum for a tetramethylene derivative **15** and then is slightly decreased for compound **16** which has a hexamethylene linker.

5

Table 6. Activities of Quinolin-4-amines **11–18** and Quaternary Derivatives **17–19**



10


No.	R	n	EC ₅₀ (nM)
11	H	3	4400
12	<i>p</i> -tolyl	3	11.5
13	2-naphthyl	3	11.0
14	H	4	316
15	2-naphthyl	4	4.0
16	2-naphthyl	6	5.9
17			>10,000
18	2-naphthyl		7600
19	H		>10,000

15

Methylation of the terminal dimethylamino group of 15 rendered the resultant trimethylammonium derivative 17 completely inactive. An additional methylation of 17 at the ring nitrogen atom restored some activity in the resultant dication 18. However, a dication 19 devoid of the naphthyl group was completely inactive again. Our additional studies (not shown) consistently indicated that compound 1 and its numerous analogs with a severely sterically hindered amino function at the terminus of the side chain showed comparable activities. Accordingly, the lack of activity of 17 and 19 cannot be explained in terms of an increased bulkiness of the trimethylammonium substituent in comparison to that of the dimethylamino group.

Analogues of 1 containing an alkyl group at N4 of the quinoline were inactive as well (not shown; see, however, 26 in Table 5). On the other hand, quinolines that contain groups capable of a hydrogen bonding interaction, such as a urethane in 20 or a hydroxy group in 21–24 (Table 7) show some activity. The activity of 25, the side chain of which contains additional amino functions, is greater.

Table 7. Activities of *N*-Substituted Quinolin-4-amines




No.	R	EC ₅₀ (nM)
20	(CH ₂) ₂ NHCO ₂ Bu- <i>t</i>	570
21	CH(Me)(CH ₂) ₃ C(Me) ₂ OH	1150
22	CH(Me)CH(Ph)OH (erythro)	1000
23	(CH ₂) ₂ O(CH ₂) ₂ OH	330
24	(CH ₂) ₆ OH	170
25	(CH ₂) ₃ N(CH ₂ CH ₂) ₂ N(CH ₂) ₃ NHC(O)(C H ₂) ₃ OH	25.1

This activity pattern is retained in a series of substituted 4-anilinoquinolines 26-32 (Table 8). Compound 32 has a relatively basic quinoline ($pK_a = 6.9$, a calculated value), contains hydroxy and amino functions, and is the most potent antagonist of immunostimulatory CpG-ODN's thus known.

5

Table 8. Activities of 4-(Substituted Anilino)quinolines

No.	R ¹	R ²	EC ₅₀
(nM)			
26	H	<i>n</i> -C ₄ H ₉	
>10,000			
27	H	CH ₂ CH ₂ OH	510
28	H	OH	320
29	morpholinomethyl	OH	10.0
30	piperidinomethyl	OH	1.5
31	pyrrolidinomethyl	OH	1.2
32	<i>N</i> -methylpiperazinomethyl	OH	0.24



In order to better understand the requirements for activity, eleven 2-(2-naphthyl)quinolin-4-amines of Tables 7 and 8, all containing a hydroxy group at the side chain and occasionally substituted with additional amino groups, were subjected to correlation analysis by using calculated connectivity indices. These parameters, depending on their order, encode the length and the size of a substituent and also the type and position of branching. The lipophilicity, length, and branching of hydrocarbon substituents are well characterized by the set of $^3X_p^v$ indices (Mokrosz *et al.*, 1992).

A statistically significant QSAR correlation for the eleven substituents, all containing free (non-protonated) amino groups at the side chain, was obtained (eq 1).

$$\log (1/EC_{50}) = 0.534 (\pm 0.08) ^3X_p^v + 5.600 (\pm 0.27)$$

(1)

$n = 11$ (C4 groups of 21-25 and 27-32), $r = 0.934$, $s = 0.49$, $F_{0.01} =$

61.09

A slightly inferior correlation was obtained for the same set of substituents with protonated amino groups excepting the N4 atom at the quinoline, which is non-basic due to the strong conjugation effect (not shown).

These results and the experimental results obtained with quaternary compounds 17-19 indicate that the basic amino group ($pK_a > 8$) at the side chain of an active agent is not protonated in the compound-bioreceptor complex. On the other hand, the observed pK_a correlation (Table 5) and the activity of the N1-quaternized quinoline 18 (The inactive compounds 17, 19 and others ($EC_{50} > 10,000$ nM) were assayed at the highest concentration of 10,000 nM and showed no effect on the cell culture. The EC_{50} value for 18 is 7600 ± 1200 nM.) are consistent with protonation of the relatively less basic ring nitrogen ($pK_a < 8$). It can be suggested that the free

amino group is located in a lipophilic microenvironment of the bioactive complex and is involved in hydrogen-bonding interaction with a hydrogen donor of the receptor.

Since nucleic acids are involved and several compounds of this study have been shown previously to strongly stabilize a triple-helix DNA structure, it was hypothesized initially that stabilization of the triple helix complex is part of the mechanism. However, while the isomeric compounds **1** and **3** are antagonists of the CpG-induced effect and compound **1** is one of the best triplex DNA stabilizing agents known to date, the sterically hindered isomer **3** does not promote triplex DNA formation and does not bind to the triple DNA structure (Strekowski *et al.*, 1996). While compound **1** binds weakly to duplex DNA, compound **3** does not bind at all. The two compounds do not stabilize a quadruplex DNA as well. In addition, compound **1** binds weakly to RNA (Wilson *et al.*, 1993). Thus, it can be suggested that the mechanism of quinoline antagonists does not involve binding to CpG-ODN.

4-Quinolinamines that inhibit CpG-ODN-induced effects are weak bases. Weak bases tend to partition into acidified vesicles within cells (de Duve *et al.*, 1974). Chloroquine has been well studied in this regard. It concentrates in lysosomes and collapses the pH gradient (Ohkuma and Poole, 1981), leading to the hypothesis that organelle acidification is required for intracellular recognition of CpG-ODN (Yi *et al.*, 1998). However, this hypothesis is not consistent with the activity of a diquatery derivative **18**, which is not a base. In addition, a number of independent experiments consistently indicated that the active compounds do not inhibit the action of CpG-ODN by interfering with the acidification of vesicles, nor with the uptake or subcellular distribution of CpG-ODN (see below). In summary, it seems most likely that these agents inhibit the detector system for CpG-ODN by a specific mechanism, not by a bulk effect.

EXAMPLE 2: Materials and Methods

A. Cell Culture

WEHI 231 murine B-lymphoma cells were maintained at log phase in medium as previously described (Macfarlane and Manzel, 1998).

B. Analogs

The structures of the analogs the inventors used are shown in FIG 1. The pyrimidine derivatives 215, 227, 228 and 231 (Strekowski *et al.*, 1991) and 2-naphthylquinolines 91, 267, 350 and 352 (Strekowski *et al.*, 1992; Streckowski *et al.*, 1994) were prepared by using general synthetic methodologies developed for similar compounds. Synthetic details will be reported elsewhere. All compounds were at least 98% pure as indicated by elemental analysis and analysis of their proton NMR spectra. The dimeric compounds 322 and 329 have been described (Ismail *et al.*, 1996). The other analogs have been previously reported (Macfarlane and Manzel, 1998) and were kindly supplied by Dr Jill Johnson from the National Cancer Institute, or purchased. The analogs were dissolved in dimethyl sulfoxide at a concentration of 10mM, and thereafter diluted in culture medium to the desired concentration.

C. Efficacy of Analogs

The efficacy of analogs as inhibitors of CpG-ODN-induced responses was determined using [³H]-thymidine uptake by WEHI-231 cells treated for 24 hr with anti-surface-IgM (which kills the cells by apoptosis), ODN1760 (which protects against this cell killing), and a range of concentrations of an analog (which reverses the protection by CpG-ODN) as previously described in detail (Macfarlane and Manzel, 1998).

D. Cellular Uptake of Analogs

Cellular uptake of analogs was estimated by fluorescence. WEHI-231 cells (5×10^6 /ml) were incubated in RPMI 1640 without phenol red (Gibco BRL, Gaithersburg, MD) with the indicated concentration of analog for 30 min. The cells were pelleted by centrifugation, and the fluorescence of the supernatant was measured

after dilution with the appropriate buffer. The cell pellet was extracted with 1 ml Catrimox- 14 (Iowa Biotechnology Corp., Coralville, LA), and the insoluble nucleic acids were removed by centrifugation ($16,000 \times g$, 10 min). The resulting extract was diluted with the appropriate buffer, and its fluorescence was measured. Preliminary experiments revealed that the Catrimox-14 extract contained essentially all of the cell-associated analog. The fluorescence intensity and spectra were determined with a Farrand Manual Spectrofluorometer at three pH's (3.4, 7.4, 10.0). Subsequent measurements of fluorescence were performed at pH giving the highest yield (Table I). Fluorescence intensity was converted to concentration using standard curves prepared by diluting each analog (0.1-5 μM) in Catrimox- 14- and RPMI-containing buffer.

E. Uptake of CpG-ODN

WEHI 231 cells were incubated with 5 $\mu\text{g/ml}$ Texas Red labeled ODN 1760. After incubation with the additions indicated, the cells were centrifuged and resuspended three times into Hanks buffered salt solution at 37° to remove surface bound ODN, and then resuspended and fixed in 2% paraformaldehyde for analysis in the University of-Iowa Flow Cytometry Facility. 10,000 events were recorded using a Coulter EPICS 753 at 488/600nm, gating out dead cells. The mean cellular fluorescence was recorded.

F. pH of CpG-ODN-Containing Cells

The pH of CpG-ODN within cells was estimated by taking advantage of the fact that acidification suppresses the fluorescence of fluorescein (Tonkinson and Stein, 1994; Ohkuma and Poole, 1978). WEHI-231 cells ($1 \times 10^6/\text{ml}$) were incubated with fluorescein-labeled ODN 1760 (5 $\mu\text{g/ml}$, 2 hr), and washed as described above. The fluorescence of the live cells was determined by flow cytometry immediately after the addition of analogs or monensin (10 μM). The pH of the ODN containing compartment was determined from the increase in the fluorescence that occurred when monensin was added, assuming that monensin equilibrates this compartment with the medium (pH 7.4). The inventors calibrated this assay using a standard curve

of fluorescence of fluorescein-labeled ODN 1760 at a range of pH's, which procedure yielded a pK_a of 6.98 in 25mM Tris acetate/150mM NaCl buffer.

G. Confocal Microscopy

WEHI 231 cells were incubated with 5 µg/ml Texas Redlabeled ODN 1760 plus additions as indicated. The cells were washed three times to remove surface bound ODN, centrifuged onto glass slides using a cytocentrifuge, and immediately fixed with 2% paraformaldehyde. The cells were photographed with a BioRad (Richmond, CA) MRC-1024 confocal laser scanning imaging system.

H. Confocal Microscopy of Live Cells

Preliminary studies established that paraformaldehyde does not prevent the leaching of quinacrine (Duve *et al.*, 1974) or other analogs from the cells. The distribution of these compounds was therefore observed in live cells. WEHI 231 cells were incubated with 50 nM quinacrine for 30 min. Cells were then wet mounted on a Fisher brand charged slide, covered with a coverslip, and photographed live as above. For comparison, cells were also incubated with 5 µg/ml Texas Red-labeled ODN 1760 for 2 hr, washed, resuspended in RPMI 1640 without phenol red and wet mounted live.

I. NOESY of ODN Interaction with Phospholipid

Preparation of vesicles was based on the procedures of Bammel *et al.* (Bammel *et al.*, 1986). Compound 91 in solid form was added to the vesicle suspension that was then stirred for 12-15 hr at 35° C. This addition procedure avoids changing the solvent composition and follows the post-membrane-preparation extrinsic ligand addition practice used in functional preparations. The compound 91 content in the DMPC vesicles was limited to 8 mol percent to avoid phase separation in the bilayer.

Phase-sensitive two-dimensional NOESY spectra of sonicated DMPC vesicles to which compound 91 was bound were obtained using a Varian Unity Plus spectrometer operating at a proton frequency of 500 MHz. The pulse sequence

developed by States *et al.* (States *et al.*, 1982) was used for data acquisition. The probe temperature was set at 35°C for all studies. An optimum spectral width of 4773.3 Hz was determined for the initial NOESY study and maintained for all subsequent acquisitions. An optimum balance of sensitivity and resolution was obtained by using 256 t1 increments each consisting of 2048 data points that were signal-averaged over 128 scans with a relaxation delay of 2 s and 215 ms acquisition time while mixing times of 50 and 150 ms were employed. Very similar NOESY spectra were obtained using both of the aforementioned mixing times. In order to minimize the possible appearance of artifactual cross peaks due to spin diffusion, NOESY spectra obtained with a 50 ms mixing time only are considered and presented herein. Suppression of the residual HOD signal was accomplished by saturation with the receiver at a power setting of 24 Hz [1/(4 PW90)]. Data processing was performed using the program VNMR supplied by Varian Associates, Inc. The free induction decays were zero filled to 2K data points in the t1 dimension. The data presented as NOESY spectra were analyzed using a sinebell window function shifted by 60°. Linear prediction (Gray, 1990) was used to obtain the first two points in the FID's and to extend the number of increments from 256 to 512 in the F1 dimension. Using the algorithm of Brown (Brown, 1995), base line correction was applied to both dimensions after fourier transformation of data in the t2 and t1 dimensions. A fourth order polynomial was usually employed for these corrections. The two-dimensional contour plots are shown in pure absorption phase.

J. DNA Thermal Melting

Thermal melting studies were conducted at 260 nm with a Cary spectrophotometer interfaced to a microcomputer. A thermistor fixed into a reference cuvette was used to monitor the temperature. The DNA oligomer was added to 1 mL of buffer (MES with 150 mM KCl, pH 4.4) in 1 cm path length reduced-volume quartz cells. Studies were generally conducted at a concentration of 5×10^{-5} M bases. Studies with chloroquine 15 and quinacrine 17 were at a ratio of 0.3 moles of compound per base.

K. CpG oligonucleotides

CpG-ODN 1760, synthesized with a phosphorothioate backbone, has the sequence:

5'-ATAATCGACGIIICAAGCAAG3'

5 The fluorescent ODNs were ODN 1760 with either fluorescein or Texas Red linked to the 5'-terminus. The oligodeoxynucleotides were purchased from Genosys (The Woodlands, TX). 1,2-dimyristoyl-*sn*-glycero-3-phosphocholine (DMPC) was obtained in powder form from Avanti Polar Lipids, Alabaster, AL, USA. The lipid was prepared synthetically and was at least 99.9 percent pure. ACS reagent grade KCl and dibasic potassium phosphate were obtained from the Fisher Scientific Co. D₂O at 10 99.9 percent purity was obtained from either Cambridge Isotope Laboratories, Woburn, MA or Isotec, Inc., Miamisburgh, OH. Monensin, chloroquine, quinine, and quinacrine were purchased from Sigma Chemical Co. (St. Louis, MO). Hydroxychloroquine was purchased from Copley Pharmaceutical, Inc. (Canton, MA). 15 Charged microscope slides were purchased from Fisher Scientific (Pittsburgh, PA).

EXAMPLE 3: Cellular Uptake of Analogs

The ability of numerous novel analogs of chloroquine and quinacrine to block the activity of CpG-ODN was examined. From the library a number of compounds 20 were selected that are fluorescent (including quinacrine) and that have a range of potency as inhibitors of CpG-ODN (FIG. 2, Table 10). Standard curves of the fluorescence intensity of each analog in cell culture medium and in the cell extract buffer were prepared, measurements being taken at the pH listed in the table. To examine the uptake of the analogs by cells, WEHI 231 cells were incubated with a 25 range of concentrations of the analogs, and measured the concentration of the analog remaining in the supernatant and the amount of analog in the extract of a cell pellet using the standard curves.

FIG. 3 shows some of the results. The cells incorporated substantial amounts 30 of the analogs (often more than half of the added reagent). The volume of each cell is about 1.0 nl (by measuring the volume of a cell pellet). Using this value, it was

estimated that the cells accumulated analogs to a concentration several hundred fold higher than in the medium (Table 10). Similar results have been reported for chloroquine e.g.(Ohkuma and Poole, 1981; Fitch *et al.*, 1974). The inventors found a linear relationship between uptake of each of the compounds and its free
5 concentration. No evidence was found to suggest that the uptake process was saturable in the low micromolar range. It was found that (non-fluorescent) chloroquine (up to 10 μ M) did not inhibit the uptake of other analogs, which also suggests that the transport mechanism is not saturable. This result with WEHI 231 cells differs from the uptake of chloroquine by red blood cells infected with malaria
10 parasites, which others have reported to be saturable and inhibitable by analogs, suggesting that infected red cells have a high affinity transport system for chloroquine (Fitch *et al.*, 1974).

The uptake of the fluorescent analogs was rapid, reaching equilibrium in less
15 than two minutes. The efflux analogs from cells when they are diluted into fresh medium was also rapid (FIG. 4).

The inventors found that uptake of quinacrine was partially inhibited by the addition of monensin, an ionophore that collapses pH gradients across biological
20 membranes. Monensin also induced the loss of quinacrine 17 from cells preloaded with the analog. Both effects were half maximal at a monensin concentration of about 1 μ M (FIG. 5). Similar results were obtained with other analogs. This result is consistent with equilibration of analogs (as weak bases) into acidic vesicles.

Table 10: Analogs Used

Our Number, source	Fluorescence vs quinacrine, pH	Name	50% Efficacy, nM	Cellular uptake ratio	Molecules/cell at 50% efficacy
329, d	1.2, b	Ism 16	14.5	147	1.28
24, a	0.7, n	10591-T/2	30.2	161	2.92
17, a	1.0, b	Quinacrine	10.2	601	3.69
25, b	0.7, a	10593-V/2	19.1	438	5.03
322, d	0.6, b	Ism 9	105.0	86	5.47
29, b	1.7, n	30020-N/2	57.5	164	5.69
30, b	1.7, n	74602-V/4	52.4	299	9.44
27, b	3.3, a	30018-L/1	60.3	583	21.1
19, a	0.8, pH2.0	Quinine	23000	24	331.1
267, c	2.2, a	OZ-66	>3000	362	>654
227, c	1.4, n	MC-241	>3000	63	>115
228, c	1.3, n	MC-132	>3000	27	>49
215, c	0.7, a	RW-15	>3000	70	>126
231, c	0.3, n	DH-77	>3000	29	>52
91, c	ND	LS-8	9.5	ND	ND
350, c	ND	OZ-123	5.9	ND	ND
352, c	ND	MHQ-6	16.3	ND	ND

Table 10 (Continued)

Our Number, source	Fluorescence vs quinacrine, pH	Name	50% Efficacy, nM	Cellular uptake ratio	Molecules /cell at 50% efficacy
15, a	Non-fluorescent	Chloroquine	110.0.0	ND	ND
16, a	Non-fluorescent	Hydroxy - chloroquine	407	ND	ND

The sources were a=Sigma Chemical Co., b=National Cancer Institute, c=Synthesized by LS, d=Synthesized by FMI. The second column is the fluorescence intensity (uncorrected) compared to quinacrine, recorded in a buffer at one of three pH's: a=3.8, n=7.4, b=10.0. The fluorescence of quinine was recorded at pH 2.0. The Efficacy is the concentration required for 50% blockage of the ability of CpG-ODN to rescue WEHI 231 cells from anti-surface IgM-induced cell death. The cellular uptake ratio is the concentration of compound in live WEHI 231 cells (assumed to have a volume of 1 nl) to the concentration in the medium at equilibrium. The molecules at 50% efficacy is the calculated number of molecules in the cell required for 50% inhibition of CpG-ODN effect. ND, not determined.

EXAMPLE 3: Comparison of Uptake and Activity

The ability of analogs to inhibit an immunostimulatory effect of CpG-ODN was examined. This assay measures the reversal of the protection by CpG-ODN against surface IgM induced killing of WEHI 231 cells. The activity of analogs was compared to their uptake, exposing a poor correlation between these two parameters (Table 10). At half maximal efficacy, cells accumulated a few million molecules of the most active analogs.

EXAMPLE 4: Effect on CpG-ODN Uptake

The effect of chloroquine and non-fluorescent analogs on the uptake of Texas Red-labeled CpG-ODN 1760 was investigated. This derivatized ODN is active as an immune stimulator. Flow cytometry measures cell-associated fluorescence because signals derived from dead cells (which take up ODN avidly, (Tonkinson and Stein, 1994)) are easily gated out of the analysis. The analogs did not significantly alter the uptake of the fluorescent ODN (FIG. 6).

Using confocal microscopy, the inventors determined that the fluorescence of Texas Red-labeled CpG-ODN 1760 is confined to numerous small organelles distributed throughout the cytoplasm with a slight predominance in the perinuclear region and Golgi apparatus. Chloroquine 15 did not influence the subcellular distribution of internalized CpG-ODN.

EXAMPLE 5: Effect on CpG-ODN pH

The fluorescence of fluorescein is suppressed at acid pH. Cells preloaded with fluorescein-labeled CpG-ODN were used to determine the pH of cell-associated ODN using flow cytometry. The addition of monensin immediately before flow cytometry results in increased fluorescence, attributable to the equilibration of the pH of the ODN with the pH of the medium and the diluent used in flow cytometry.

A standard curve was implemented to relate fluorescence with pH, enabling the pH of the cell compartment occupied by ODN to be determined to be about 6.4, a value very close to that published by Tonkinson *et al.* using a similar method (Tonkinson and Stein, 1994). Using this method, the inventors examined the effects of (non-fluorescent) analogs on the pH of the compartment. The effect of 1 μ M chloroquine (a concentration that completely suppresses CpG-ODN responses) was an increase in the pH of CpG-ODN-containing compartment by less than 0.01 pH units. No analog increased the pH substantially at the concentration required for activity, although they did collapse the pH at higher concentration.

EXAMPLE 6: Subcellular Localization of Analogs

Next, the subcellular localization of fluorescent analogs was examined using confocal microscopy. Unable to find a method of fixation that immobilized the analogs within the cells, the inventors incubated the cells with the analogs and examined them live. A consistent difference between the distribution of CpG-ODN and of fluorescent analogs was revealed: the analogs occupy organelles that were larger and more peripherally located within the cell. However, it is not clear that quinacrine does not also enter the compartment occupied by CpG-ODN, because photographs of cells double-labeled with Texas-Red labeled CpG-ODN and quinacrine 17 did not unambiguously reveal a population of ODN-containing vesicles that also did not stain with quinacrine.

EXAMPLE 7: Interaction between Analogs and ODN

Structural nucleic acids with duplex or single-stranded folded conformations generally give sigmoid thermal melting curves characteristic of thermal unfolding of the structures. This unfolding results in a change in the UV spectrum. In other studies, the CpG ODN 1760 gave a very small continuous increase in absorbance at 260 nm as a function of temperature, characteristic of some base unstacking in single-stranded DNA, and did not show any transition curve characteristic of a base-paired structure. Addition of chloroquine 15 or quinacrine 17 to the CpG-ODN did not cause any significant change in the thermal melting curve of the free CpG-ODN. It can, thus, be concluded that under these conditions, these compounds bind poorly, if at all, to the CpG-ODN.

EXAMPLE 8: Interaction between Analogs and Lipids

The structures of compound 91 and DMPC are provided in FIG. 9; the lipid NMR resonance assignments that are listed in Table 11 are keyed to the numbered groups on the DMPC structure in this figure. The assignments in Table 11 are derived in part from the work of Ellena *et al.*, 1987 with additional assignments by the inventors.

Table 11: Proton resonance Chemical Shift Assignments for Compound 91 and DMPC in Unilamellar Vesicles

Proton Number	Group	Chemical Shift, in D ₂ O, (ppm)
14	-CH ₃	0.87
13	-CH ₂ -	1.17
12	-CH ₂ -	1.25
4-11	-(CH ₂) ₄₋₁₁ -	1.29
3	-CH ₂ -	1.56
2	-CH ₂ -	2.33
15	CHOCO-	5.26 – 5.31
16	H ₂ COCO-	4.43
16	H ₂ COPO- (glyceride)	3.99
17	-OPOCH ₂ - (choline)	4.27
18	-CH ₂ N-	3.56 – 3.63
19	N(CH ₃) ₃	3.12 – 3.19
Compound 91	Naphthalene, quinoline	7 - 9

5

A typical spectrum obtained at a pH of 7. is shown in FIG. 9 and is referenced to the terminal methyl resonance of the DMPC acyl chain (0.87 ppm); this assignment is relative to an external TMS standard. NOESY cross peaks between the compound 91 quinoline and naphthalene ring protons and those of the DMPC fatty acid unresolved methylenes (1.3 ppm on the F1 axis, Nos. 4-11; FIG. 9A) are present. Additional cross peaks are observed between the compound 91 aromatic ring protons and those of the lipid glycerol backbone methylenes (3.99 ppm, No. 16) and to those of the choline N-CH₂ moiety (3.5 ppm, No. 18). Very weak, marginal cross peaks are observed between the compound 91 aromatic system protons and those of the choline N(CH₃)₃ moiety (3.2 ppm, No. 19) and to those of the fatty acid terminal methyl group at 0.87 ppm on the F1 axis (No. 14, FIG. 9A). The NOESY spectrum obtained

10

15

at a pD of 4.7 is qualitatively similar to that obtained at a pD of 7.4. In the aliphatic region, the cross peak at 4.25 ppm on the F2 axis is assigned to the interaction of the compound 350 aliphatic chain methylene groups with those of the OPO-CH₂ methylene (No. 17) (FIG. 9B).

5

A location model that is consistent with the NOESY data suggests that the charged N(CH₃)₂ moiety be located no deeper in the membrane bilayer than the onset of the hydrocarbon region - near the ester linkages in the DMPC lipids. The NOESY results indicate that the aliphatic chain of compound 91 is located near the interface of the choline head group and the glycerol backbone of the lipid. Such a location would allow for favorable electrostatic interactions between the protonated N(CH₃)₂ group and the negatively charged DMPC phosphate moiety.

10

The unambiguous cross peaks between the aromatic ring protons of compound 91 and the unresolved DMPC fatty acid methylene protons (1.3 ppm on the F1 axis) indicates that a portion of the compound 91 molecules penetrate the bilayer to at least the level of the fourth methylene group on the fatty acid chain and perhaps below this level.

15

Molecular modeling studies using SYBYL (Tripos Associates) of a DMPC bilayer with the lipids in the all *trans* conformation and with compound 91 in an energy minimized extended conformation with the protonated N(CH₃)₂ nitrogen located near the phosphate group will allow an NOE cross peak between protons on the naphthalene ring and the No. 8 methylene protons with a 5 Å NOE cutoff distance criterion.

20

25

The lack of cross peaks between the terminal methyl group and the aromatic protons does not preclude the eventual translocation of compound 91 through the bilayer and the accumulation of the drug candidate in the vesicle internal volume as has been observed in whole cell studies based on similar quinolines tagged with fluorescent labels as described in a separate section of this communication. The NMR results suggest in this instance that the steady state population of compound 91

30

molecules near the middle portion of the bilayer, which would include the roughly 0.5 mol percent of compound 91 in neutral form, is below the detection threshold for the NOESY experiment in the DMPC vesicle system.

5 **EXAMPLE 9: Preparation of 2-(Substituted phenyl)-4-quinolinamines**

***N*-[3-(Dimethylamino)propyl]-2-[4-(*N*-methylpiperazino)phenyl]quinolin-4-amine (44).** A ketimine 36 derived from 33 and 34 was cyclized to a quinoline 38 in the presence of potassium *tert*-butoxide by using a procedure for the preparation of similar compounds (Strekowski *et al.*, 1997). Hydrolysis of 38 in the presence of acid followed by treatment of the resultant hydroxyquinoline 40 with POCl₃ gave a chloroquinoline 42. The general procedures are also described in the same reference. The selective displacement of the chlorine atom in 42 was accomplished by treatment with *N,N*-dimethyl-1,3-propanediamine in the presence of a catalytic amount of SnCl₄ (4h, 130°C). Stirring of the resultant product (1 mmol) with lithium *N*-methylpiperizide (20 mmol) in anhydrous THF (50 mL) at 23 °C for 20h followed by quenching with water gave compound 44 that was purified by silica gel chromatography eluting with EtOAc/NEt₃ (17:3): yield 61%, mp of the hydrobromide salt (x3HBr) 274-276 °C.

20 ***N*-[3-(Dimethylamino)propyl]-2-[2-[[2-(dimethylamino)ethyl]amino]-phenyl]quinolin-4-amine (45).** A ketimine 37 was obtained from 33 and 35 as indicated above. Its cyclization to 39, hydrolysis of 39 to 41, and the synthesis of 43 were carried out as described above as well. Following the treatment of 43 with *N,N*-dimethyl-1,3-propanediamine the resultant product was allowed to react with excess lithium 2-(dimethylamino)ethylamide in THF at 23 °C for 20h. Product 45 was purified by silica gel chromatography eluting with EtOAc/Et₃N (25:1): yield 76%, a solid.

30 ***N*-(3-Morpholinopropyl)-2-[4-[[2-pyridyl)methylamino]methyl]phenyl]-quinoline (49).** Synthesis of 4-chloro-2-(4-tolyl)quinoline (46) was conducted by using the chemistry described above. The treatment of 46 (5 mmol) with *N*-bromosuccinimide (5 mmol) in the presence of benzoyl peroxide (100 mg) in CCl₄

(50 mL) under reflux for 6h gave **47**: yield 63%, mp 116-117°C (from hexanes). A mixture of **47** (0.6 mmol) and 2-(aminomethyl)pyridine (2 mL) was stirred at 23°C for 6h and then treated with water (20 mL). Extraction with ethyl acetate followed by drying of the extract and concentration gave compound **48** that was used for the reaction with 3-morpholinopropylamine without any purification. A mixture of **48**, the amine (1 mL), and a catalytic amount of SnCl₄ was heated under reflux for 4h and then quenched with water (20 mL). Following a standard workup, crude product **49** was purified by silica gel chromatography eluting with EtOAc/MeOH/Et₃N (10:1:3): yield 58%, mp 140-142 °C.

EXAMPLE 10: Preparation of Substituted 4-Quinolinamines and 9-Aminoacridines

Compounds have been numbered in this example. This numbering refers only to the compounds found in this example and in Scheme 4 shown in FIG. 11, which corresponds to this example.

Materials and Methods

Melting points (Pyrex capillary) are uncorrected. ¹H NMR (300 MHz) and ¹³C NMR (75.4 MHz) spectra were taken with TMS as an internal reference. Proton-proton coupling constants smaller than 2 Hz are not reported. Ketimine **1** was obtained as reported previously (Strekowski *et al.*, 1992).

Reaction of Ketimine 1 with Potassium *tert*-Butoxide.

A mixture of **1** (10 g, 32 mmol) and *t*-BuOK (17 g, 150 mmol) in anhydrous THF (500 mL) was heated under reflux under a nitrogen atmosphere for 1 h. Cooling to 23° C was followed by quenching with water (6.0 mL) and filtration. Concentration of the filtrate on a rotary evaporator to 50 mL followed by dilution with hexanes (40 mL) gave a precipitate of **5**, which was filtered off and crystallized from methanol. Flash chromatography of the THF/hexanes solution (silica gel; hexanes/Et₃N, 95:5) gave an analytically pure sample of compound **4**, which was additionally purified by crystallization from hexanes. Crude compound **4** was used for the subsequent reaction.

4-(*tert*-butoxy)-2-(2-naphthyl)quinoline (4): 1.06 g (51%); mp 76-78°C; ¹H NMR (CDCl₃) δ 1.64 (s, 9 H), 7.19 (s, 1 H), 7.41 (t, J = 8 Hz, 1 H), 7.46 (m, 3 H), 7.63 (t, J = 8 Hz, 1 H), 7.83 (m, 1 H), 7.93 (m, 2 H), 8.06 (d, J = 8 Hz, 1 H), 8.14 (d, J = 8 Hz, 1 H), 8.19 (d, J = 8 Hz, 1 H), 8.45 (s, 1 H); ¹³C NMR (CDCl₃) δ 28.9, 80.7, 105.1, 122.6, 123.2, 125.2, 125.3, 126.3, 1256.6, 126.9, 127.7, 128.5, 128.8, 12802, 129.8, 133.4, 133.7, 137.8, 149.8, 158.1, 160.1, Anal. Calcd for C₂₃H₂₁NO: C, 84.39; H, 6.46; N, 4.28. Found C, 84-39; H, 6.59; N, 4.45.

2-(2-naphthyl)quinolin-4(1H)-one (5): yield 1.3 G (15%); mp 290-292°C (from MeOH); ¹H NMR (DMSO-*d*₆) δ 3.13 (br s, 1 H), 6.48 (s, 1 H), 7.35 (t, J = 8 Hz, 1H), 7.63 (m, 2 H), 7.69 (t, J = 8 Hz, 1H), 7.81 (d, J = 8 Hz, 1 H), 7.93 (d, J = 8 Hz, 1 H), 8.03 (m, 1 H), 8.11 (m, 2 H), 8.46 (s, 1 H); ¹³C NMR (DMSO-*d*₆) δ 107.8, 118.7, 123.3, 124.5, 124.8, 124.9, 127.0, 127.2, 127.5, 127.7, 128.6, 128.7, 131.5, 131.9, 132.6, 133.6, 140.6, 149.9, 176.9; IR (KBr) ν 3250, 1630, 1595, 1547, 1508 cm⁻¹. Anal. Calcd for C₁₉H₁₃NO: C, 84.11; H, 4.83; N, 5.16. Found: C, 83.80; H, 4.69; N, 5.14.

4-Hydroxy-2-(2-naphthyl)quinolinium *p*-toluenesulfonate (6). A solution of crude compound 4 (6.5 g, 20 mmol) and *p*-toluenesulfonic acid (6.0 g, 30 mmol) in THF (150 mL) was heated under reflux for 4 h and then cooled to 0°C, the precipitate of 6 was filtered off and crystallized from methanol: yield 8.0 g (93%); mp 223-225°C; ¹H NMR (DMSO-*d*₆) δ 2.28 (s, 3), 7.10 (d, J = 8 Hz, 2 H), 7.28 (s, 1 H), 7.50 (d, J = 8 Hz, 2 H), 7.71 (m, 3 H), 8.0-8.215 (m, 6 H), 8.33 (d, J = 8 Hz, 1 H), 8.61 (s, 1 H). Anal. Calcd for C₂₆H₂₁NO₄S: C, 70.40; H, 4.77; N, 3.16. Found: C, 70.38; H, 4.73; N, 3.11.

4-Chloro-2-(2-naphthyl)quinoline (7). A mixture of salt 6 (7.0 g, 15.7 mmol), phosphorus pentachloride (3.2 g, 15.7 mmol), and phosphorus oxychloride (40 mL) was heated under reflux for 1 h, then cooled, and poured onto ice. The mixture was neutralized with a saturated solution of sodium bicarbonate, and the resultant

precipitate of crude product 7 was filtered, washed with water, and dried (50°C/10 mmHg). Purification involved treatment with hot ethyl acetate (180 mL), filtration from an insoluble yellow solid, then concentration of the solution, and crystallization of the residue from hexanes: yield 3.98 g (86%); mp 114-116°C; ¹H NMR (CDCl₃) δ 7.48 (m, 2 H), 7.58 (t, J = 8 Hz, 1 H), 7.74 (t, J = 8 Hz, 1 H), 7.84 (m, 1 H), 7.94 (m, 2 H), 8.08 (s, 1 H), 8.17 (m, 2 H), 8.28 (d, J = 8 Hz, 1 H), 8.53 (s, 1 H); ¹³C NMR (CDCl₃) δ 119.5, 124.3, 125.0, 125.7, 126.8, 127.3, 127.6 (two signals), 128.0, 129.0, 129.2, 130.4, 130.9, 133.7, 134.4, 136.1, 143.5, 149.4, 157.3. Anal. Calcd for C₁₉H₁₂ClN: C, 78.75; H, 4.17; N, 4.71. Found: 78.75; H, 4.11; N, 4.65.

The treatment of quinoline 5 as described above furnished product 7 in a similar yield.

2-(2-Naphthyl)-4-[(p-tolysulfonyl)oxy]quinoline (8). A solution of salt 6 (4.3 g, 10 mmol) in pyridine (60 mL) was cooled to 0°C and treated slowly with *p*-toluensesulfonyl chloride (3.4 g, 18 mmol) at such a rate that the temperature did not rise above 5°C. After being stirred for an additional 3 h at 5°C, the mixture was diluted with dichloromethane (40 mL) and poured into cold water (200 mL). The organic layer was washed in succession with 1 N hydrochloric acid, a solution of sodium bicarbonate, and water, then dried over sodium sulfate, and concentrated on a rotary evaporator. Chromatography on silica gel eluting with hexanes/*tert*-butyl methyl ether (1:1) followed by crystallization from *tert*-butyl methyl ether gave 3.5 g (79%) of 8: mp 128-130°C; ¹H NMR (DMSO-*d*₆) δ 2.43 (s, 3 H), 7.51 (d, J = 8 Hz, 2 H), 7.62 (m, 3 H), 7.86 (d, J = 8 Hz, 2 H), 7.96 (m, 3 H), 8.02 (m, 1 H), 8.13 (m, 3 H), 8.34 (d, J = 8 Hz, 1 H), 8.65 (s, 1 H); ¹³C NMR (CDCl₃) δ 21.7, 110.5, 121.2, 121.4h, 124.7, 126.5, 126.9, 127.0, 127.2, 127.7, 128.6, 128.7, 128.8, 129.5, 130.1, 130.6, 132.2, 133.3, 134.0, 136., 146.1, 150.1, 153.8, 157.8. Anal. Calcd for C₂₆H₁₉NO₃S: C, 73.39; H, 4.50; N, 3.29. Found: , 73.12; H, 4.32; N, 3.27.

N-[3-[4-[3-(4-Hydroxybutyramido)propyl]piperazino]-propyl]-2-(2-naphthyl)quinolin-4-amine (9). A mixture of 7 (5.0 g, 17 mmol), 1,4-bis(3-

aminopropyl)piperazine (58 mL, 282, mmol), and anhydrous tin tetrachloride (0.8 mL, 6.6 mmol) was stirred and heated to 130 °C under a nitrogen atmosphere for 3.5 h. Removal of the excess of the amine (100 °C/0.2 mmHg) was followed by addition of γ -butyrolactone (3 mL, 39 mmol) and stirring of the mixture at 80°C for an additional
5 2.5 h. After cooling to 23°C, water (10 mL) was added and the resultant solution was extracted with ethyl acetate (6 x 30 mL). The extract was dried over magnesium sulfate and concentrated, and the residue was crystallized from anhydrous ethyl acetate: yield 5.1 g (56%); mp 177-179 °C; ¹H NMR (DMSO-*d*₆) δ 1.5-3.5 (m, 26 H), 4.42 (t, J = 5 Hz, 1 H, exchangeable with D₂O), 7.14 (s, 1 H), 7.38 (br, 1 H, exchangeable with D₂O), 7.42 (t, J = 8 Hz, 1 H), 7.55 (m, 2 H), 7.65 (t, J = 8 Hz, 1 H),
10 7.73 (br, 1 H, exchangeable with D₂O), 7.91 (d, J = 8 Hz, 1 H), 7.96 (m, 1 H), 8.02 (d, J = 8 Hz, 1 H), 8.09 (m, 1 H), 8.21 (d, J = 8 Hz, 1 H), 8.39 (d, J = 8 Hz, 1 H), 8.70 (s, 1 H); ¹³C NMR (DMSO-*d*₆) δ 24.9, 26.3, 28.6, 32.1, 36.8, 41.3, 52.7, 53.0, 55.5, 56.1, 60.3, 95.1, 118.0, 121.4, 123.7, 125.0, 126.1, 126.2, 126.4, 127.4, 127.7, 128.5, 129.1,
15 129.3, 132.9, 133.2, 137.5, 148.2, 150.9, 156.4, 171.9. Anal. Calcd for C₃₃H₄₁N₅O₂: C, 73.43; H, 7.65; N, 12.98. Found: C, 73.08; H, 7.71; N, 12.89.

A hemihydrate 9•½ H₂O was obtained by crystallization of crude 9 from wet ethyl acetate, mp 181-182 °C. Anal. Calcd for 9•½ H₂O: C, 72.22; H, 7.71; N, 12.76.
20 Found: C, 72.19; H, 7.76; N, 12.72.

Discussion

Several synthetic approaches to the title compound and analogs have been evaluated. This compound is a practical precursor to *N*-substituted 2-(2-naphthyl)quinolin-4-amines, the triple-helix DNA specific intercalators.
25

The triple-helix structure of nucleic acids is formed by binding a single strand of DNA in the major groove of duplex DNA. The interaction is highly base sequence specific but is quite unstable under normal physiological conditions. Various
30 biotechnology applications have led to an increased interest in stabilization of the triplex DNA form. For example, the formation of a stable triplex structure between a

short oligonucleotide and a specific sequence in a long duplex DNA can be used to inhibit expression of the specific gene. Alternatively, with an appropriate cleaving group attached to the third-strand oligomer, highly specific cleavage of DNA can be achieved.

5

One approach to enhance triplex stability is to design compounds that bind strongly and specifically to triplex DNA but weakly to duplex DNA. Another strategy is to tether such compounds to the triplex-forming oligonucleotide, so that the triplex structure can be stabilized efficiently by intramolecular interactions.

10

Several DNA intercalators are known to interact nonselectively with triple and double DNA structures or stabilize the triple helix relative to the corresponding duplex with various selectivities. Compound 2 and other *N*⁴-substituted 2-(2-naphthyl)quinolin-4-amines are far superior in their triplex stabilization ability and the triplex/duplex binding selectivity than any other triplex intercalators reported to date. These unfused bi-aromatic derivatives bind to and stabilize strongly and selectively T-AT triplets of the triple-helix DNA in the presence of duplex DNA of any sequence.

15

In this study critically examine several synthetic approaches to such triplex DNA intercalators. The preparation of a standard intercalator 2 and the synthesis of compound 9 with a terminal hydroxy group for the attachment to the 5'-end of an oligonucleotide by using phosphoramidite chemistry serve as examples. A similar strategy can be used for the synthesis of analogs of 9 containing a terminal 1,2-diol functionality for linking to the 3'-end of an oligonucleotide.

20

25

Quinoline 2 has been obtained previously by two methods, namely, (i) lithium 2-(dimethylamino)-ethylamide mediated cyclization of ketimine 1 derived from 2-(trifluoromethyl)aniline and acetophenone and (ii) nucleophilic displacement of fluoride in 4-fluoroquinoline 3 with *N,N*-dimethylethylenediamine. Unfortunately, the short and efficient route (i) is not applicable to the preparation of other quinolines substituted with a primary alkylamino function at position 4.

30

The attractiveness of the second method (ii) as a general route to 4-(substituted alkylamino) quinolines is severely hampered by the low yield of 4-fluoroquinoline 3 obtained by the reaction of 2-(trifluoromethyl)aniline with the lithium enolate of acetophenone and a tedious purification that requires several consecutive chromatographic separations. Numerous attempts to optimize the synthesis of 3 did not succeed. The 33% yield of 3 obtained on a 200-mg scale decreased to 5-15% for the reactions conducted on a 2-g scale under a variety of experimental conditions. These experiments included the published conditions and reactions conducted in different solvents (ether, hexanes, and ether/hexanes, in the presence and absence of hexamethylphosphoramide) with varying ratios of the reagents and at varying temperatures.

Inventors focused their attention on 2-arylquinolin-4(1H)-ones that could be transformed into the corresponding quinolines substituted with other nucleofugal groups at position 4. A method described by Staskun failed to produce the desired 2-(2-naphthyl)quinolin-4(1H)-one (5). The successful synthetic route to 5 involves potassium *tert*-butoxide mediated cyclization of ketimine 1. This reaction furnished a mixture of a 4-*tert*-butoxyquinoline 4 and quinolin-4(1H)-one 5 in 51% and 15% yields, respectively. The *tert*-butoxy derivative 4 was efficiently converted into 6 (93% yield), a *p*-toluenesulfonate salt of 5, by treatment with *p*-toluene-sulfonic acid. The subsequent reaction of either 5 or 6 with phosphorus oxychloride afforded 4-chloro-2-(2-naphthyl)-quinoline (7) in 75% yield. The use of phosphorus oxychloride improved the efficiency of this reaction to 86%. Larger amounts of phosphorus pentachloride did not increase the yield of 7 but used potential safety hazards during workup. Substitution of *p*-toluenesulfonyl chloride for POCl₃ furnished 2-(2-naphthyl)-4-[(*p*-tolylsulfonyl)oxy]quinoline (8). Unfortunately, the treatment of 8 with a primary amine did to cause displacement of the sulfonate group but resulted in cleavage of the S-O bond and gave quinoline 5 quantitatively. Initial attempts at displacing chloride from 7 by primary amines also did not succeed. Thus, in contrast to the reactivity of its fluoro analog 3, the 4-chloroquinoline 7 was inert in attempted reactions with amines at 120°C. After 6 h at 150°C, the starting material 7 was still present and a large number of products had formed, as observed by TLC analysis.

The nucleophilic displacement of chloride easily took place, however, in the presence of a catalytic amount of tin tetrachloride. Such a treatment of 7 with *N,N*-dimethylethylenediamine gave the known intercalator 2. In the synthesis of the desired triplex DNA intercalator 9, compound 7 was treated with an excess of 1,4-bis(3-aminopropyl)piperazine in the presence of SnCl₄ followed by a reaction of the resultant intermediate mono-substituted amine, without purification, with γ -butyrolactone. The efficiencies of the reactions conducted with 0.5 g of 7 or scaled up to 5 g were virtually identical. The yield of the two-step transformation of 7 to 9 was 56%, and the overall yield of 9 from 1 was 27%.

In summary, a practical method is described for the synthesis of 4-chloro-2-(2-naphthyl)quinoline (7) and shown that compound 7 is a good substrate for the preparation of triple-helix DNA intercalators. Products, such as 9, that are compatible with phosphoramidite chemistry can be used in the preparation of triple-helix DNA intercalator-oligonucleotide conjugates.

EXAMPLE 11: Preparation of bis-4-quinolinamines and bis-9-aminoacridines

Materials and Methods

General Procedure. A mixture of 4-chloro-2-(2-naphthyl)quinoline (300 mg, 1.0 mmol), an α,ω -diamine (H-R-H, 0.05 mmol), and a catalytic amount of SnCl₄ (30 μ L) was heated in a Parr bomb to 140 °C for 3.5 h. *N,N*-Bis[3-[2-(2-naphthyl)quinolin-4-amino]propyl]methylamine was crystallized as a hydrobromine salt, and the purification of 1,4-Bis[3-[2-(2-naphthyl)quinolin-4-amino]propyl]piperazine and *N,N*-Bis[2-(2-naphthyl)quinolin-4-yl]]4,9-dioxa-1,12-dodecanediamine was conducted by silica gel chromatography of free bases eluting with AcOEt followed by crystallization of the free bases.

***N,N*-Bis[3-[2-(2-naphthyl)quinolin-4-amino]propyl]methylaniline:** yield 52% of $C_{45}H_{41}N_5 \cdot 3HBr \cdot H_2O$; mp 257-262°C; 1H NMR δ 2.29 (m, 4H), 2.82 (s, 3H), 3.28 (m, 4H), 3.82 (m, 4H), 7.27 (s, 2H), 7.64 (m, 6H), 7.92 (t, $J = 8$ Hz, 2H), 8.02 (d, $J = 8$ Hz, 2H), 8.15 (t, $J = 8$ Hz, 4H), 8.21 (t, $J = 8$ Hz, 4H), 8.69 (d, $J = 8$ Hz, 2H), 8.80 (s, 2H), 9.50 (br, exchangeable with D_2O), 11.10 (br, exchangeable with D_2O). Anal. Calcd for $C_{45}H_{41}N_5 \cdot 3HBr \cdot H_2O$: C, 59.40; H, 4.88; N, 7.70. Found: C, 59.70; H, 5.12; N, 7.59.

1,4-Bis[3-[2-(2-naphthyl)quinolin-4-amino]propyl]piperazine: yield 60% of the free base; mp 192-198 °C; 1H NMR δ 1.90 (m, 4H), 2.48 (m, 4H), 3.35 (s, 8H), 3.52 (m, 4H), 7.15 (s, 2H), 7.44 (t, $J = 8$ Hz, 2H), .50 (br, exchangeable with D_2O), 7.55 (m, 4H), 7.66 (t, $J = 8$ Hz, 2H), 7.92 (d, $J = 8$ Hz, 2H), 7.96 (m, 2eH), 8.04 (d, $J = 8$ Hz, 2H), 8.09 (m, 2H), 8.24 (d, $J = 8$ Hz, 2H), 8.38 (d, $J = 8$ Hz, 2H), 8.17 (s, 2H). Anal. Calcd for $C_{48}H_{46}N_6 \cdot H_2O$: C, 79.52; H, 6.67; N, 11.60. Found: C, 79.39; H, 7.04; N, 11.70.

***N,N*-Bis[2-(2-naphthyl)quinolin-4-yl][4,9-dioxa-1,12-dodecanediamine:** yield 40% of the free base; mp 115-117 °C; 1H NMR δ 1.44 (m, 4H), 1.93 (m, 4H), 3.31 (m, 4H), 3.42 (m, 4H), 3.58 (m, 4H), 7.10 (s, 2H), 7.53 (t, $J = 8$ Hz, 2H), 7.57 (m, 4H), 7.78 (t, $J = 8$ Hz, 2H), 7.96 (m, 2H), 8.06 (m, 6H), 8.22 (m, 6H), 8.22 (d, $J = 8$ Hz, 2H), 8.26 (br, exchangeable with D_2O), 8.41 (d, $J = 8$ Hz, 2H), 8.67 (s, 2H). Anal. Calcd for $C_{48}H_{46}N_4O_2 \cdot H_2O$: C, 79.09; H, 6.64; N, 7.69. Found: C, 78.70; H, 6.69; N, 7.41.

Bis[3-[2-(2-naphthyl)quinolin-4-amino]propyl] Succinate. A solution of 4-chloro-(2-naphthyl) quinoline (420 mg, 1.4 mmol) in 3-aminopropanol (1.0 mL) was heated to 130°C for 1.5 h under a nitrogen atmosphere. Treatment with water (2 mL) was followed by extraction of the mixture with AcOEt, drying of the extract with $MgSO_4$, and concentration to give a residue of crude *N*-(3-hydroxypropyl)-2-(2-naphthyl)quinolin-4-amine. This compound was purified by silica gel chromatography eluting with $CH_2Cl_2/MeOH$ (5:1) followed by crystallization of its

hydrobromide salt; yield 82% of $C_{22}H_{20}N_2O \cdot HBr \cdot H_2O$; mp 230-235 °C. Anal. Calcd for $C_{22}H_{20}N_2O \cdot HBr \cdot H_2O$: C, 61.82; H, 5.43; N, 6.55. Found: C, 61.98; H, 5.28; N, 6.49.

5 A mixture of $C_{22}H_{20}N_2O \cdot HBr \cdot H_2O$ (530 mg, 1.25 mmol), $SOCl_2$ (1.0 mL), and benzene (5 mL) was heated under reflux for 1.5 h. Following concentration on a rotary evaporator, the oily residue was treated with AcOEt (25 mL) and an aqueous solution of $NaHCO_3$ (10%, 10 mL), and the mixture was stirred briefly. The organic layer was separated, dried ($MgSO_4$), and concentrated. The residue of crude *N*-(3-chloropropyl)-2-(2-naphthyl)quinolin-4-amine was purified by silica gel chromatography eluting with hexanes/ether (1:2) and then crystallized; yield 75%; mp 131-135 °C. HRMS. Calcd for $C_{22}H_{19}^{35}ClN_2$: m/z 346.1254. Found m/z 346.1240.

15 *N*-(3-chloropropyl)-2-(2-naphthyl)quinolin-4-amine (208 mg, 0.6 mmol), succinic acid (30 mg, 0.3 mmol), and DBU (90 μ L, 0.6 mmol) in anhydrous DMF (1.0 mL) was heated to 90 °C for 6 h under a nitrogen atmosphere. Following concentration on a rotary evaporator, the residue of bis[3-[2-(2-naphthyl)quinolin-4-amino]propyl] succinate was subjected to silica gel chromatography eluting with CH_2Cl_2/i -PrOH (5:1). Then it was transformed into a hydrobromide salt, and the salt was crystallized; yield 30% of $C_{48}H_{42}N_4O_4 \cdot 3HBr \cdot 2.5H_2O$; mp 210-216 °C; 1H NMR δ 2.10 (m, 4H), 2.52 (s, 4H), 4.04 (m, 4H), 4.15 (m, 4H), 7.65 (m, 10H), 7.79 (d, J = 8 Hz, 2H), 7.98 (m, 4H), 8.07 (m, 4H), 8.14 (d, J = 8 Hz, 2H), 8.30 (s, 2H), 8.52 (d, J = 8 Hz, 2H). Anal. Calcd for $C_{48}H_{42}N_4O_4 \cdot 3HBr \cdot 2.5H_2O$: C, 56.13; H, 4.87; N, 5.45, Found: C, 56.03; H, 4.62; N, 5.52.

25

Quinolin-4-amines. A mixture of 4-chloro-(2-naphthyl) quinoline (600 mg, 2.1 mmol), *N,N*-bis(3-aminopropyl)methylamine or 1,3-propanediamine (3 mL), and $SnCl_4$ (50 μ L) was heated to 140°C for 4 h under a nitrogen atmosphere. Removal of excess amine under reduced pressure was followed by treatment of the residue with hydrobromic acid and then crystallization of the resultant salt.

30

N-[3-[*N*-3-Aminopropyl)methylamino]propyl]quinolin-4-amine: yield 64% of $C_{26}H_{30}N_4 \cdot 3HBr$; mp 232-235 °C; 1H NMR δ 2.02 (m, 2H), 2.23 (m, 2H), 2.84 (s, 3H), 2.92 (m, 2H), 3.28 (m, 4H), 3.83 (m, 2H), 7.13 (s, 1H), 7.16 (m, 2H), 7.30 (t, $J = 8$ Hz, 1H), 7.91 (br, exchangeable with D_2O), 8.02 (t, $J = 8$ Hz, 1H), 8.11 (d, $J = 8$ Hz, 1H), 8.20 (m, 4H), 8.71 (d, $J = 8$ Hz, 1H), 8.81e (s, 1H), 9.44 (br, exchangeable with D_2O), 9.82 (br, exchangeable with D_2O). Anal. Calcd for $C_{26}H_{30}N_4 \cdot 3HBr$: C, 35.43; H, 4.11; N, 6.36. Found: C, 35.53; H, 3.99; N, 6.12.

N-(3-Aminopropyl)-2-(2-naphthyl)quinolin-4-amine: yield 61% of $C_{22}H_{21}N_3 \cdot 2HBr \cdot 1.5H_2O$; mp 280-284°C; 1H NMR δ 2.13 (m, 2H), 3.03 (m, 2H), 3.24 (br, exchangeable with D_2O), 3.85 (m, 2H), 7.27 (s, 1H), 7.70 (m, 3H), 7.95 (br, exchangeable with D_2O), 7.99 (t, $J = 8$ Hz, 1H), 8.07 (d, $J = 8$ Hz, 1H), 8.18 (m, 3H), 8.24 (t, $J = 8$ Hz, 1H), 8.71 (d, $J = 8$ Hz, 1H), 8.79 (s, 1H), 9.37 (br, exchangeable with D_2O). Anal. Calcd for $C_{22}H_{21}N_3 \cdot 2HBr \cdot 1.5H_2O$: C, 51.17; H, 5.09; N, 8.14. Found: C, 51.19; H, 4.85; N, 7.96.

1,4-Bis[3[(3-carboxypropionamido)propyl]]piperazine. A solution of 1,4-bis(3-aminopropyl)piperazine (1.0 mL, 4.8 mmol) and succinic anhydride (1.0 g, 10.0 mmol) in anhydrous DMF (20 mL) was heated to 70°C for 1 h under a nitrogen atmosphere. The resultant precipitate was filtered, washed with EtOH, and crystallized from DMF: yield 57%; mp 179-180°C; 1H NMR (D_2O) δ 1.87 (m, 4H), 2.51 (m, 8H), 2.93 (m, 4H), 3.24 (m, 12H). Anal. Calcd for $C_{18}H_{32}N_4O_6$: C, 54.00; H, 8.05; N, 14.00. Found: C, 53.99; H, 8.20; N, 13.98.

A General Procedure for the Preparation of the Remaining Three Compounds. A mixture of succinic acid or 1,4-bis[3[(3-carboxypropionamido)propyl]]piperazine (0.3 mmol), triethylamine (0.25 mL, 1.8 mmol), and BOP (225 mg, 0.6 mmol) in anhydrous DMF (4 mL) was stirred at 23°C for 15 min before treatment with a salt $C_{26}H_{30}N_4 \cdot 3HBr$ or $C_{22}H_{21}N_3 \cdot 2HBr \cdot 1.5H_2O$ (0.6 mmol). The mixture was stirred at 23 °C for 24 h and then filtered. The solution

was concentrated on a rotary evaporator and a residue of a crude product was transformed into a hydrobromide salt, and the salt crystallized.

***N,N*-Bis[3-[*N*-[3-[2-(2-naphthyl)quinolin-4-amino]propyl]methylamino]-propyl]succinamide** (from *N*-[3-[*N*-3-aminopropyl]methylamino]propyl]quinolin-4-amine and succinic acid): yield 73% of $C_{56}H_{62}N_8O_2 \cdot 5HBr \cdot 5H_2O$; mp 232-240°C; 1H NMR δ 1.81 (m, 4H), 2.19 (m, 4H), 2.31 (s, 3H), 2.78 (s, 4H), 3.09 (m, 8H), 3.27 (m, 4H), 3.81 (m, 4H), 7.27 (s, 2H), 7.71 (m, 6H), 8.01 (m, 4H), 8.10 (m, 2H), 8.16 (m, 4H), 8.23 (d, $J = 8$ Hz, 2H), 8.62 (d, $J = 8$ Hz, 2H), 8.74 (s, 2H), 9.30 (br, exchangeable with D_2O), 9.55 (br, exchangeable with D_2O). Anal. Calcd for $C_{56}H_{62}N_8O_2 \cdot 5HBr \cdot 5H_2O$: C, 49.02; H, 5.24; N, 8.19. Found: C, 48.89; H, 5.35; N, 8.55.

1,4-Bis[3-[[*N*-[3-[2-(2-naphthyl)quinolin-4-amino]propyl]succinamoyl]amino]-propyl]piperazine (from *N*-(3-aminopropyl)-2-(2-naphthyl)quinolin-4-amine and 1,4-bis[3[(3-carboxypropionamido)propyl]]piperazine): yield 47% of $C_{62}H_{70}N_{10}O_4 \cdot 4HBr$; mp 290-296°C; 1H NMR δ 1.86 (m, 8H), 2.64 (s, 8H), 2.7-3.6 (m, 24H), 7.21 (s, 2H), 7.71 (m, 6H), 7.87 (br, exchangeable with D_2O), 8.01 (t, $J = 8$ Hz, 2H), 8.15 (m, 10H), 8.62 (d, $J = 8$ Hz, 2H), 8.71 (s, 2H). Anal. Calcd for $C_{62}H_{70}N_{10}O_4 \cdot 4HBr$: C, 44.78; H, 4.60; N, 8.42. Found: C, 44.69; H, 4.86; N, 8.24.

1,4-Bis[3-[[*N*-[3-[*N*-[2-[2-(2-naphthyl)quinolin-4-amino]propyl]methylamino]propyl]succinamoyl]amino]propyl]piperazine (from *N*-[3-[*N*-3-aminopropyl]methylamino]propyl]quinolin-4-amine and 1,4-bis[3[(3-carboxypropionamido)propyl]]piperazine): yield 46% of $C_{70}H_{88}N_{12}O_4 \cdot 9HBr \cdot 2H_2O$; mp 234-240°C; 1H NMR δ 1.89 (m, 8H), 2.08 (m, 4H), 2.32 (m, 4H), 2.56 (m, 8H), 2.65 (s, 8H), 2.82 (s, 6H), 2.94 (m, 4H), 3.11 (m, 4H), 3.34 (m, 4H), 3.87 (m, 8H), 7.30 (s, 2H), 7.71 (m, 6H), 7.93 (br, exchangeable with D_2O), 8.01 (t, $J = 8$ Hz, 2H), 8.08 (d, $J = 8$ Hz, 2H), 8.21 (m, 8H), 8.74 (d, $J = 8$ Hz, 2H), 8.81 (s, 2H), 9.38 (br, exchangeable with D_2O), 9.93 (br, exchangeable with D_2O). Anal. Calcd for

$C_{70}H_{88}N_{12}O_4 \cdot 9HBr \cdot 2H_2O$: C, 43.64; H, 5.28; N, 8.73. Found: C, 43.81; H, 5.43; N, 8.46.

5

All of the compositions and methods disclosed and claimed herein can be made and executed without undue experimentation in light of the present disclosure. While the compositions and methods of this invention have been described in terms of preferred embodiments, it will be apparent to those of skill in the art that variations may be applied to the compositions and methods and in the steps or in the sequence of steps of the method described herein without departing from the concept, spirit and scope of the invention. More specifically, it will be apparent that certain agents that are both chemically and physiologically related may be substituted for the agents described herein while the same or similar results would be achieved. All such similar substitutes and modifications apparent to those skilled in the art are deemed to be within the spirit, scope and concept of the invention as defined by the appended claims.

10

15

REFERENCES

The following references, to the extent that they provide exemplary procedural or other details supplementary to those set forth herein, are specifically incorporated herein by reference.

U.S. Patent No. 4,016,100

U.S. Patent No. 4,089,801

U.S. Patent No. 4,234,871

U.S. Patent No. 4,485,054

Ballas, Rasmussen, Krieg, "Induction of NK activity in murine and human cells by CpG motifs in oligodeoxynucleotides and bacterial DNA," *J. Immunology*, 157:1840-1845, 1996.

Bammel, Brand, Germon, Smith, "Interaction of the extrinsic potential-sensitive molecular probe diS-C3- (5) with pigeon heart mitochondria under equilibrium and time-resolved conditions," *Arch. Biochem. Biophys.*, 244:67-84, 1986.

Bangham *et al.*, *J. Mol. Biol.* 13:238-252, 1965.

Brown, "Fully automated baseline correction of 1D and 2D NMR spectra using bernstein polynomials," *Magn. Reson.*, Series A 114:268-270, 1995.

Deamer and Uster, "Liposome Preparation: Methods and Mechanisms," *In: Liposomes*, M. Ostro (Ed.), 1983

de Duve, C.; de Barsey, T.; Poole, B.; Trouet, A.; Tulkens, P.; van Hoof, F. *Biochem. Pharm.*, 23, 1974.

Drug Carriers In Biology and Medicine, G. Gregoriadis ed., pp. 287-341, 1979.

Duve, Barsey, Poole, Trouet, Tulkens, van Hoof, "Lysosomotropic agents," *Biochem. Pharm.*, 23, 1974.

Ellena, Dominey, Archer, Xu, Cafiso, "Localization of hydrophobic ions in phospholipid bilayers using ¹H nuclear Overhauser effect spectroscopy," *Biochem.*, 26:4584-4592, 1987.

Fitch, Yunis, Chevli, Gonzalez, "High affinity accumulation of chloroquine by mouse erythrocytes infected with *Plasmodium berghei*," *J. Clin. Invest.*, 54:24-33, 1974.

Fox, "Mechanism of action of hydroxychloroquine as an antirheumatic drug," *Sem. Arthritis Rheumatism*, 23:82-91, 1993.

Ghosh and Bachhawat, Targeting of Liposomes to Hepatocytes. In: *Liver Diseases, Targeted Diagnosis and Therapy Using Specific Receptors and Ligands*. Wu et al., eds., Marcel Dekker, New York, pp. 87-104, 1991.

Gray, "Linear prediction guide," *Magn. Moments*, 7:30-33, 1990.

Ismail, Dascombe, Carr, North, "An exploration of the structure-activity relationships of 4-aminoquinolines: novel antimalarials with activity *in-vivo*," *J. Pharm. Pharmacol.*, 48:841-850, 1996.

10 Krieg et al., "Lymphocyte activation by CpG dinucleotide motifs in prokaryotic DNA," *Trends Microbiol.*, 4(2):73-6, 1996.

Krieg, Matson, Fisher, "Oligodeoxynucleotide modifications determine the magnitude of B cell stimulation by CpG motifs," *Antisense Nucleic Acid Drug Dev.*, 6(2):133-9, 1996.

15 Krieg, Yi, Matson, Waldschmidt, Bishop, Teasdale, Koretzky, Klinman, "CpG motifs in bacterial DNA trigger direct B-cell activation," *Nature*, 374:546-549, 1995.

MacFarlane and Manzel, "Antagonism of immunostimulatory CpG-oligodeoxynucleotides by quinacrine, chloroquine and structurally related compounds," *J. Immunol.*, 160:1122-1131, 1998.

20 MacFarlane, Manzel, Krieg, "Unmethylated CpG-containing oligodeoxynucleotides inhibit apoptosis in WEHI 231 B-lymphocytes induced by several agents: evidence for blockade at a distal signaling step," *Immunology*, 91:586-593, 1997.

25 Mokrosz, J. L.; Duszynska, B.; Strekowski, L. *Pharmazie*; and references cited therein, 47:538, 1992.

Ohkuma and Poole, "Cytoplasmic vacuolation of mouse peritoneal macrophages and the uptake into lysosomes of weakly basic substances," *J. Cell. Biol.*, 90:656-664, 1981.

30 Ohkuma and Poole, "Fluorescence probe measurements of the intralysosomal pH in living cells and the perturbation of pH by various agents," *Proc. Natl. Acad. Sci. USA*, 75:3327-3331, 1978.

- Remington's Pharmaceutical Sciences, 15th ed., Mack Publishing Company, Easton, PA, 1980.
- States, Haberkorn, Rubin, "A two-dimensional nuclear overhauser experiment with pure absorption phase in four quadrants," *Magn. Reson.*, 48:286-292, 1982.
- 5 Strekowski, Kiselyov, Hojjat, "The o-amino-trifluoromethyl functionality as a novel synthon for 4-fluoroquinolines," *J. Org. Chem.*, 59:5886-5890, 1994.
- Strekowski, L.; Gulevich, Y.; Baranowski, T. C.; Parker, A. N.; Kiselyov, A. S.; Lin, S.-Y.; Tanious, F. A.; Wilson, W. D. *J. Med. Chem.* 39:3980, 1996.
- Strekowski, L.; Janda, L.; Patterson, S. E.; Nguyen, J. *Tetrahedron*, 52:3273, 1996.
- 10 Strekowski, L.; Zegrocka, O.; Windham, C.; Czarny, A. *Org. Process Res. Dev.*, 1:384, 1997.
- Strekowski, Patterson, Janda, Wydra, Harden, Lipowska, Cegl, "Further studies on the cyclization of aromatic azaomethines ortho-substituted with a trifluoromethyl group: Synthesis of 2,4-di- or 2,3,4-trisubstituted quinolines," *Org. Chem.*, 57,
- 15 1992.
- Strekowski, Wilson, Mokrosz, Mokrosz, Harden, Tanious, Wydra, Crow, Jr., "Quantitative structure-activity relationship analysis of cation-substituted polyaromatic compounds as potentiators (amplifiers) of bleomycin-mediated degradation of DNA," *J. Med. Chem.*, 34:580-588, 1991.
- 20 Szoka and Papahadjopoulos, *Proc. Natl. Acad. Sci. USA*, 75: 4194-4198, 1978.
- Tonkinson and Stein, "Patterns of intracellular compartmentalization, trafficking and acidification of 5'-fluorescein labeled phosphodiester and phosphorothioate oligodeoxynucleotides in HL60 cells," *Nucleic Acid Res.*, 22:4268-4275, 1994.
- Wallace, "Antimalarial agents and lupus," *Rheumatic Disease Clinics of Portly*
- 25 *America*, 20:243-263, 1994.
- Wawrzynczak & Thorpe "Methods for Preparing Immunotoxins: Effect of the Linkage on Activity and Stability", In: Immunoconjugates: Antibody Conjugates in Radioimaging and Therapy of Cancer, Vogel (ed.), New York, Oxford University Press, pp. 28-55, 1987.
- 30 Wilson, 1998. "DNA and RNA intercalators," In: *DNA and aspects of molecular biology*, Kool (ed.), Vol. 7, 1998.


Wilson, W. D.; Ratmeyer, L.; Zhao, M.; Strekowski, L.; Boykin, D. *Biochemistry*, 32:4098, 1993.

Yi, Hornbeck, Lafrenz, Krieg, "CpG DNA rescue of murine B lymphoma cells from anti-IgM induced growth arrest and programmed cell death is associated with expression of *c-myc*, *c-myb*, *myn*, and *bcl-2*," *J. Immunol.*, 157:4918-4925, 1996.

Yi, Tuetken, Redford, Waldschmidt, Kirsch, Krieg, "CpG motifs in bacterial DNA activate leukocytes through the pH-dependent generation of reactive oxygen species," *J. Immunol.*, 160:4755-4761, 1998.

CLAIMS:

1. A method of inhibiting immunostimulation in a subject, the method comprising administering an effective amount of a substituted 4-quinolinamine composition to said subject, the 4-quinolinamine composition comprising a compound having the structural formula A



wherein

RA is hydrogen atom or a lower alkyl group;

RB is a substituted or unsubstituted alkyl, alkenyl or alkynyl secondary or tertiary amine;

R2 is a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, an substituted or unsubstituted anthracyl group, a substituted or unsubstituted phenanthryl group or a substituted or unsubstituted styryl group;

R3 is a hydrogen atom;

R5 is a hydrogen atom;

R6 is a hydrogen atom or a halogen atom;


R7 is a hydrogen atom or a halogen atom; and

R8 is a hydrogen atom,

and pharmaceutically acceptable salts thereof.

2. The method of claim 1, wherein said R2 phenyl, naphthyl, anthracyl, styryl or phenanthryl group substitution is further defined as one or more substitutions selected from the group consisting of an alkyl group, an alkoxy group, an alkoxyalkyl group, an ester group, an alkylamino group, a dialkylamino group, a cyclic amino group, a halogen atom and any combination thereof.

3. The method of claim 2, wherein said cyclic amino group is a piperazino group, a piperidino group, a pyrrolidino group, an imidazolyl group, pyridyl group, or a morpholino group.
4. The method of claim 1, wherein said R₈ alkyl substitution is selected from the group consisting of a cyclic amino group, an alkylamino group, a dialkylamino, furyl, phenyl, thienyl, and any combination thereof.
5. The method of claim 4, wherein said cyclic amino group is a piperazino group, a piperidino group, a pyrrolidino group, an imidazolyl group, a pyridyl group, or a morpholino group.
6. A method of inhibiting immunostimulation in a subject, the method comprising administering an effective amount of a substituted 4-quinolinamine composition to said subject, the 4-quinolinamine composition comprising a compound having the structural formula B



wherein the phenyl group can be substituted at R₉, R₁₀, R₁₁, R₁₂ and R₁₃, wherein said substituted phenyl group comprises one or more substitutions selected from the group consisting of an alkyl group, an alkoxy group, a hydroxy group, an alkylaminoalkyl group, an alkoxyalkyl group, an ester group, an alkylamino group, a dialkylamino group, a cyclic amino group, a furan group, a thiophene group, a halogen atom or any combination thereof;

R₂ is a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted anthracyl group, a substituted or unsubstituted styryl group or a substituted or unsubstituted phenanthryl group;

R₃ is a hydrogen atom;

R₅ is a hydrogen atom;

R₆ is a hydrogen atom or a halogen atom;

R₇ is a hydrogen atom, a halogen atom or an alkyl halogen atom;

R₈ is a hydrogen atom,

and pharmaceutically acceptable salts thereof.


7. The method of claim 6, wherein said cyclic amino group is a piperazino group, a piperidino group, a pyrrolidino group, an imidazolino group or a morpholino group.

8. The method of claim 7, wherein said piperidino group is 4-alkyl piperazino.

9. The method of claim 6, wherein the R₂ phenyl, naphthyl, anthracyl, styryl or phenanthryl group substitution is further defined as one or more substitutions selected from the group consisting of an alkyl group, an alkoxy group, a hydroxy group, an alkoxyalkyl group, an ester group, an alkylamino group, an dialkylamino group, a cyclic amino group, a halogen atom or any combination thereof.

10. The method of claim 9, wherein said cyclic amino group is a piperazino group, a piperidino group, a pyrrolidino group, an imidazolino group or a morpholino group.

11. A method of inhibiting immunostimulation in a subject, the method comprising administering an effective amount of a substituted 4-quinolinamine composition to said subject, the 4-quinolinamine composition comprising a compound having the structural formula C



wherein

the phenyl group can be unsubstituted or substituted at R₉, R₁₀, R₁₁, R₁₂ and R₁₃, wherein said substituted phenyl group comprises one or more substitutions selected from the group consisting of an alkyl group, an alkoxy group, an alkoxyalkyl group, an alkylaminoalkyl group, a hydroxy group, an ester group, an alkylamino group, an dialkylamino group, a cyclic amino group, a furan group, a thiophene group, a halogen atom, or any combination thereof;

n is 0 to 4 CH₂;

R₂ is a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted anthracyl group, a substituted or unsubstituted styryl group or a substituted or unsubstituted phenanthryl group;

R₃ is a hydrogen atom;

R₅ is a hydrogen atom;

R₆ is a hydrogen atom or a halogen atom;

R₇ is a hydrogen atom, a halogen atom or an alkyl halogen atom;

R₈ is a hydrogen atom,


and pharmaceutically acceptable salts thereof.

12. The method of claim 11, wherein said cyclic amino group is a piperazino group, a piperidino group, a pyrrolidino group, an imidazolino group or a morpholino group.

13. The method of claim 11, wherein the R_2 phenyl, naphthyl, anthracyl, styryl or phenanthryl group substitution is further defined as one or more substitutions selected from the group consisting of an alkyl group, an alkoxy group, hydroxy group, an alkoxyalkyl group, an ester group, an alkylamino group, an dialkylamino group, a cyclic amino group, a halogen atom or any combination thereof.

14. The method of claim 13, wherein said cyclic amino group is a piperazino group, a piperidino group, a pyrrolidino group, an imidazolino group or a morpholino group.

15. A method of inhibiting immunostimulation in a subject, the method comprising administering an effective amount of a substituted bis -4-quinolinamine composition to said subject, the bis-4-quinolinamine composition comprising a compound having the structural formula D



wherein

R_B on the first 4-quinolinamine is covalently attached to $R_{B'}$ on the second 4-quinolinamine by linker group Y;

R_2 is a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted anthracyl group, a substituted or unsubstituted styryl group or a substituted or unsubstituted phenanthryl group;

R₃ is a hydrogen atom;

R₅ is a hydrogen atom;

R₆ is a hydrogen atom or a halogen atom;

R₇ is a hydrogen atom or a halogen atom;

R₈ is a hydrogen atom;

R₂ is a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, an substituted or unsubstituted anthracyl group, a substituted or unsubstituted styryl group or a substituted or unsubstituted phenanthryl group;

R₃ is a hydrogen atom;

R₅ is a hydrogen atom;

R₆ is a hydrogen atom or a halogen atom;

R₇ is a hydrogen atom or a halogen atom; and

R₈ is a hydrogen atom,

and pharmaceutically acceptable salts thereof.

15. The method of claim 14, wherein the R₂ phenyl, naphthyl, anthracyl, styryl or phenanthryl group substitution is further defined as one or more substitutions selected from the group consisting of an alkyl group, an alkoxy group, an alkoxyalkyl group, a hydroxy group, an ester group, an alkylamino group, an dialkylamino group, a cyclic amino group, a halogen atom or any combination thereof.

16. The method of claim 15, wherein said cyclic amino group is a piperazino group, a piperidino group, a pyrrolidino group, an imidazolino group or a morpholino group.


17. The method of claim 14, wherein the R₂ phenyl, naphthyl, anthracyl, styryl or phenanthryl group substitution is further defined as one or more substitutions selected from the group consisting of an alkyl group, an alkoxy group, an alkoxyalkyl group, a hydroxy group, an ester group, an alkylamino group, an dialkylamino group, a cyclic amino group, a halogen atom or any combination thereof.

18. The method of claim 17, wherein said cyclic amino group is a piperazino group, a piperidino group, a pyrrolidino group, an imidazolino group or a morpholino group.

19. The method of claim 14, wherein said linker Y is an alkyl group, an ester group, an ether group, an alkylamino group, amido group, a cyclohexanedieyl, 1,4-piperazinedieyl or any combination thereof.

20. The method of claim 19, wherein said linker is α,ω -alkanedieyl.

21. A method of inhibiting immunostimulation in a subject, the method comprising administering an effective amount of a substituted bis-9-aminoacridine composition to said subject, the bis-9-aminoacridine composition comprising a compound having the structural formula E



wherein

R_B on the first 9-aminoacridine is covalently attached to $R_{B'}$ on the second 9-aminoacridine by linker group Y;

OR_2 is a lower alkyl group;

OR_2' is a lower alkyl group;

R_3 is a hydrogen atom or a lower alkoxy group;


R_3' is a hydrogen atom or a lower alkoxy group;

X is a halogen atom;

X' is a halogen atom,

and pharmaceutically acceptable salts thereof.

22. The method of claim 21, wherein said linker Y is an alkyl group, an ester group, an ether group, an alkylamino group, amido group, a cyclohexanedieyl, 1,4-piperazinedieyl or any combination thereof.
23. The method of claim 22, wherein said linker is α,ω -alkanedieyl.
24. A method of inhibiting immunostimulation in a subject, the method comprising administering an effective amount of a composition to said subject, the composition comprising a compound having the structural formula F



wherein

R_A is hydrogen atom or a lower alkyl group;

R_B is covalently linked to R_B' by linker Y;

R₂ is a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, an substituted or unsubstituted anthracyl group, a substituted or unsubstituted styryl group or a substituted or unsubstituted phenanthryl group;

R₃ is a hydrogen atom;

R₅ is a hydrogen atom;

R₆ is a hydrogen atom or a halogen atom;

R₇ is a hydrogen atom or a halogen atom;

R₈ is a hydrogen atom;

R_B' is linked covalently to R_B by linker Y;

OR₂' is a lower alkyl group; and

R₃' is a hydrogen atom, or a lower alkoxy group;

X is a halogen atom,

and pharmaceutically acceptable salts thereof.


25. The method of claim 24, wherein said R_2 phenyl, naphthyl, anthracyl, styryl or phenanthryl group substitution is further defined as one or more substitutions selected from the group consisting of an alkyl group, an alkoxy group, an alkoxyalkyl group, a hydroxy group, an ester group, an alkylamino group, an dialkylamino group, a cyclic amino group, a halogen atom or any combination thereof.

26. The method of claim 24, wherein said cyclic amino group is a piperazino group, a piperidino group, a pyrrolidino group, an imidazolino group or a morpholino group.

27. The method of claim 24, wherein said linker Y is an alkyl group, an ester group, an ether group, an alkylamino group, amido group, a cyclohexanedieyl, 1,4-piperazinedieyl or any combination thereof.

28. The method of claim 27, wherein said linker is α,ω -alkanedieyl.

29. A substituted 4-quinolinamine composition having the structural formula A



wherein

R_A is hydrogen atom or a lower alkyl group;

R_B is a substituted or unsubstituted alkyl, alkenyl or alkynyl secondary or tertiary amine;

R₂ is a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, an substituted or unsubstituted anthracyl group, a substituted or unsubstituted phenanthryl group or a substituted or unsubstituted styryl group;

R₃ is a hydrogen atom;

R₅ is a hydrogen atom;

R₆ is a hydrogen atom or a halogen atom;

R₇ is a hydrogen atom or a halogen atom; and

R₈ is a hydrogen atom,

and pharmaceutically acceptable salts thereof.


30. The composition of claim 29, wherein said R₂ phenyl, naphthyl, anthracyl, styryl or phenanthryl group substitution is further defined as one or more substitutions selected from the group consisting of an alkyl group, an alkoxy group, an alkoxyalkyl group, a hydroxy group, an ester group, an alkylamino group, a dialkylamino group, a cyclic amino group, a halogen atom and any combination thereof.

31. The composition of claim 30, wherein said cyclic amino group is a piperazino group, a piperidino group, a pyrrolidino group, an imidazolyl group, a pyridyl group, or a morpholino group.

32. The composition of claim 29, wherein said R_B alkyl substitution is selected from the group consisting of a cyclic amino group, furyl, thienyl, alkylamino group, dialkylamino group, phenyl, and any combination thereof.

33. The composition of claim 32, wherein said cyclic amino group is a piperazino group, a piperidino group, a pyrrolidino group, an imidazolyl group, a pyridyl group, or a morpholino group.

34. A substituted 4-quinolinamine composition having the structural formula B



wherein

the phenyl group can be substituted at R₉, R₁₀, R₁₁, R₁₂ and R₁₃, wherein said substituted phenyl group comprises one or more substitutions selected from the group consisting of an alkyl group, an alkoxy group, an alkoxyalkyl group, an alkylaminoalkyl group, an ester group, an alkylamino group, an dialkylamino group, a cyclic amino group, a furan group, a thiophene group, a halogen atom or any combination thereof;

R₂ is a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted anthracyl group, a substituted or unsubstituted styryl group or a substituted or unsubstituted phenanthryl group;

R₃ is a hydrogen atom;

R₅ is a hydrogen atom;

R₆ is a hydrogen atom or a halogen atom;

R₇ is a hydrogen atom, a halogen atom or an alkyl halogen atom;

R₈ is a hydrogen atom,

and pharmaceutically acceptable salts thereof.


35. The composition of claim 31, wherein said cyclic amino group is a piperazino group, a piperidino group, a pyrrolidino group, an imidazolino group or a morpholino group.

36. The composition of claim 35, wherein said cyclic amino group is 4-alkyl-piperazino.

37. The composition of claim 34, wherein the R_2 phenyl, naphthyl, anthracyl, styryl or phenanthryl group substitution is further defined as one or more substitutions selected from the group consisting of an alkyl group, an alkoxy group, an alkoxyalkyl group, an ester group, an alkylamino group, a hydroxy group, a dialkylamino group, a cyclic amino group, a halogen atom or any combination thereof.

38. The composition of claim 37, wherein said cyclic amino group is a piperazino group, a piperidino group, a pyrrolidino group, an imidazolino group or a morpholino group.

39. A substituted 4-quinolinamine composition having the structural formula C



wherein

the phenyl group can be unsubstituted or substituted at R₉, R₁₀, R₁₁, R₁₂ and R₁₃, wherein said substituted phenyl group comprises one or more substitutions selected from the group consisting of an alkyl group, an alkoxy group, an alkoxyalkyl group, an alkylaminoalkyl group, an ester group, an alkylamino group, an dialkylamino group, a cyclic amino group, a furan group, a thiophene group, a halogen atom, or any combination thereof;

n is 0 to 4 CH₂;

R₂ is a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted anthracyl group, a substituted or unsubstituted styryl group or a substituted or unsubstituted phenanthryl group;

R₃ is a hydrogen atom;

R₅ is a hydrogen atom;

R₆ is a hydrogen atom or a halogen atom;

R₇ is a hydrogen atom, a halogen atom or an alkyl halogen atom;

R₈ is a hydrogen atom,


and pharmaceutically acceptable salts thereof.

40. The composition of claim 39, wherein said cyclic amino group is a piperazino group, a piperidino group, a pyrrolidino group, an imidazolino group or a morpholino group.

41. The composition of claim 39, wherein the R₂ phenyl, naphthyl, anthracyl, styryl or phenanthryl group substitution is further defined as one or more substitutions selected from the group consisting of an alkyl group, an alkoxy group, an alkoxyalkyl group, an ester group, an alkylamino group, an dialkylamino group, a cyclic amino group, a halogen atom or any combination thereof.

42. The composition of claim 41, wherein said cyclic amino group is a piperazino group, a piperidino group, a pyrrolidino group, an imidazolino group or a morpholino group.

43. A substituted bis -4-quinolinamine composition having the structural formula



D

wherein

R_B on the first 4-quinolinamine is covalently attached to R_{B'} on the second 4-quinolinamine by linker group Y;

R₂ is a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, a substituted or unsubstituted anthracyl group, a substituted or unsubstituted styryl group or a substituted or unsubstituted phenanthryl group;

R₃ is a hydrogen atom;

R₅ is a hydrogen atom;

R₆ is a hydrogen atom or a halogen atom;

R₇ is a hydrogen atom or a halogen atom;

R₈ is a hydrogen atom;

R_{2'} is a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, an substituted or unsubstituted anthracyl group, a substituted or unsubstituted styryl group or a substituted or unsubstituted phenanthryl group;

R_{3'} is a hydrogen atom;

R_{5'} is a hydrogen atom;

R_{6'} is a hydrogen atom or a halogen atom;

R_{7'} is a hydrogen atom or a halogen atom;

R_{8'} is a hydrogen atom,

and pharmaceutically acceptable salts thereof.

44. The composition of claim 43, wherein the R₂ phenyl, naphthyl, anthracyl, styryl or phenanthryl group substitution is further defined as one or more substitutions selected from the group consisting of an alkyl group, an alkoxy group, an alkoxyalkyl group, an ester group, an alkylamino group, an dialkylamino group, a cyclic amino group, a halogen atom or any combination thereof.


45. The composition of claim 44, wherein said cyclic amino group is a piperazino group, a piperidino group, a pyrrolidino group, an imidazolino group or a morpholino group.

46. The composition of claim 43, wherein the R₂ phenyl, naphthyl, anthracyl, styryl or phenanthryl group substitution is further defined as one or more substitutions selected from the group consisting of an alkyl group, an alkoxy group, an ether group, an ester group, an alkylamino group, an dialkylamino group, a cyclic amino group, a halogen atom or any combination thereof.

47. The composition of claim 46, wherein said cyclic amino group is a piperazino group, a piperidino group, a pyrrolidino group, an imidazolino group or a morpholino group.

48. The composition of claim 43, wherein said linker Y is an alkyl group, an ester group, an alkoxyalkyl group, an alkylamino group, amido group, a cyclohexanedieyl, an piperazino group, 1-4, piperazinedieyl or any combination thereof.

49. A substituted bis-9-aminoacridine composition having the structural formula E



wherein

R_B on the first 9-aminoacridine is covalently attached to $R_{B'}$ on the second 9-aminoacridine by linker group Y;

OR_2 is a lower alkyl group;

$OR_{2'}$ is a lower alkyl group;

R_3 is a hydrogen atom or a lower alkoxy group;

$R_{3'}$ is a hydrogen atom or a lower alkoxy group;


X is a halogen atom;

X' is a halogen atom,

and pharmaceutically acceptable salts thereof.

50. The composition of claim 49, wherein said linker Y is an alkyl group, an ester group, an alkoxyalkyl group, an alkylamino group, amido group, a cyclohexanedieyl, an piperazino group, 1-4, piperazinedieyl or any combination thereof.

51. A 4-quinoliamine, 9-aminoacridine composition having the structural formula



F

wherein

R_A is hydrogen atom or a lower alkyl group;

R_B is covalently linked to R_B by linker Y;

R₂ is a substituted or unsubstituted phenyl group, a substituted or unsubstituted naphthyl group, an substituted or unsubstituted anthracyl group, a substituted or unsubstituted styryl group or a substituted or unsubstituted phenanthryl group;

R₃ is a hydrogen atom;

R₅ is a hydrogen atom;

R₆ is a hydrogen atom or a halogen atom;

R₇ is a hydrogen atom or a halogen atom;

R₈ is a hydrogen atom;

R_B' is linked covalently to R_B by linker Y;

OR₂' is a lower alkyl group; and

R₃' is a hydrogen atom, or a lower alkoxy group;


X is a halogen atom,

and pharmaceutically acceptable salts thereof.

52. The composition of claim 51, wherein said R₂ phenyl, naphthyl, anthracyl, styryl or phenanthryl group substitution is further defined as one or more substitutions selected from the group consisting of an alkyl group, an alkoxy group, an alkoxyalkyl group, an ester group, an alkylamino group, an dialkylamino group, a cyclic amino group, a halogen atom or any combination thereof.

53. The composition of claim 52, wherein said cyclic amino group is a piperazino group, a piperidino group, a pyrrolidino group, an imidazolino group or a morpholino group.


54. The composition of claim 51, wherein said linker Y is an alkyl group, an ester group, an alkoxyalkyl group, an alkylamino group, amido group, a cyclohexanedieyl, an piperazino group, 1-4, piperazinedieyl or any combination thereof.



9.62

Strekowski


361 OZ-148



8.94

Strekowski


359 OZ-144



8.83

Strekowski


337 OZ-83



8.74


Strekowski

348 MHQ-4



8.46


Strekowski



8.94

Strekowski


363 OZ-146



8.86

Strekowski


382 OZ-145



8.79

Strekowski


372 MHQ-17



8.69

Strekowski


371 MHQ-15



8.45

Strekowski


FIG. 1A



8.40

Strekowski


349 MHQ-5



8.27

Strekowski


177 LJ-286



8.19

Strekowski


400 MS-42



8.19


Strekowski

107 TMTN 1



8.12


Kotecka



8.30

Strekowski


350 OZ-123



8.23

Strekowski


178 SP-103



8.19

Strekowski


278 OZ-95



8.15

Strekowski


275 OZ-45



8.12

Strekowski


FIG. 1B



8.04

Strekowski


109 TMTN 11



8.04

Kotecka


318 trans



8.01

Ismail


26



7.99


NCI

197 LJ-247/1



7.96


Strekowski



8.04

Strekowski


211 M-40



8.04

Strekowski


360 OZ-153



8.00

Strekowski


94 TG-84



7.96

Strekowski


279 OZ-101



7.96

Strekowski


FIG. 1C



7.96

Strekowski


394 MHQ-42



7.94

Strekowski


179 LJ-280



7.88

Strekowski


401 MS-49



7.88


Strekowski

403 MS-41



7.86


Strekowski



7.94

Ismail


395 MHQ-46



7.91

Strekowski


396 MHQ-28



7.88

Strekowski


367 MHQ-9



7.87

Strekowski


17 Quinacrine



7.85


Sigma

FIG. 1D



7.85


Strekowski



7.85

Strekowski


323 cls



7.82

Ismail


338 OZ-85H



7.82

Strekowski


339 OZ-124



7.82

Strekowski


377 MHQ-13



7.81

Strekowski


141 MC-183



7.79

Strekowski


352 MHQ-6



7.79

Strekowski


106 GBB 1608



7.78

Kotacka

111 GBB1515



7.78

Kotacka

FIG. 1E



Strekowski



Kotecka



Strekowski




Strekowski



NCI


7/32



7.72

Kotecka


128 M-149/2



7.68

Strekowski


391 MHQ-45



7.66

Strekowski


176 TG-96




7.65

Strekowski

325 trans rr




7.65



7.69

Boykin


34



7.66

NCI


38



7.65

NCI


293 DB 431



7.65

Boykin

328 trans. (+/-)




7.65

Ismail

FIG. 1G


7/32



7.62

NCl


104 TN 121



7.60

Kotecka


269 OZ-31



7.60

Strekowski


44



7.52


NCl

105 TMTN 109



7.52


Kotecka



7.62

NCl


124 M-61



7.60

Strekowski


386 MHQ-38



7.57

Strekowski


101 TN 109



7.52

Kotecka


192 LJ-327



7.45

Strekowski


FIG. 1H



7.45

Strekowski


69



7.42

NCI


103 TN 119



7.40

Kotecka


193 LJ-297



7.38


Strekowski

32



7.37


NCI



7.43

NCI


97 AK-189



7.40

Strekowski


121 M-25



7.38

Strekowski

23



7.37

NCI

145 SP-110




7.33

Strekowski

FIG. 11


10/32



7.32

Strekowski


122 M-95



7.30

Strekowski


48



7.29

NCI


30



7.28


NCI

29



7.24


NCI



7.31

Strekowski


385 MHQ-37



7.30

Strekowski


195 LJ-267



7.29

Strekowski


174 TG-88



7.26

Strekowski


27



7.22

NCI


FIG. 1J



7.22

NCl


212 M-9A



7.19

Strekowski


80



7.15

NCl


274 OZ-44



7.14


Strekowski

56



7.12


NCl



7.20

NCl


51



7.17

NCl


196 LJ-266



7.14

Strekowski


42



7.13

NCl


93 LS-12



7.12

Strekowski


FIG. 1K



7.10

NCI


96 AK-206



7.08

Strekowski


315



7.02

Ismail


207 DH-19



7.00


Strekowski

389 MHQ-33



6.98


Strekowski



7.10

Strekowski


214 M-119



7.05

Strekowski


18 Amodiaquin



7.00

Sigma


322 trans. (+/-)



6.98

Ismail


15 Chloroquine



6.96

Sigma


FIG. 1L



6.96

Molecular Probes


100 TN 108



6.92

Kotecka


35



6.87

NCI


68



6.87


NCI

172 LJ-256/2



6.87


Strekowski



6.94

NCI


164 AK-195



6.88

Strekowski


53



6.87

NCI


163 X-37



6.87

Strekowski

61




6.86

NCI

FIG. 1M


14/32



6.85

Strekowski


263



6.85

WRAIR


50



6.83

NCl


10



6.82


NCl

12



6.80


NCl



6.85

Strekowski


9



6.84

NCl


4



6.82

NCl


198 LJ-331



6.81

Strekowski


147 LJ-250



6.79

Strekowski


FIG. 1N



6.79

Strekowski


246 DB449



6.76

Boykin


354 S-10



6.75

Strekowski


380 MHQ-25



6.71


Strekowski

194 LJ-265



6.69


Strekowski



6.77

Strekowski


64



6.75

NCI


65



6.72

NCI


1



6.70

NCI


67



6.68

NCI


FIG. 10



6.64

NCl


76



6.60

NCl


185 LJ-320



6.55


Strekowski

74



6.52


NCl



6.62

NCl


11



6.56

NCl


73



6.54

NCl


62



6.51

NCl

FIG. 1P



P.T.C. 2




FIG. 3

19/32




FIG. 4

20/32




FIG. 5




FIG. 6

22/32




FIG. 7

23/32




FIG. 8

FIG. 9A





FIG. 9B



Scheme 2




FIG. 10A

Scheme 3




FIG. 10B

27/32

Scheme 4






FIG. 11


MS-128




MS-55




MS-76




MS-98




MS-81-2



MS-12-1



MS-130



MS-95






FIG. 12A

MS-132




8.62

MS-133




8.48

MHQ-92




8.49

MS-129




8.31

MHQ-108




8.43

MS-77



8.27


MS-136



9.26


FIG. 12B

MS-140




8.26

MS-125




8.17

MHQ-88




8.22

MS-139




7.94

MHQ-71




8.17

MHQ-93




8.15

MHQ-89



8.11


MS-134



8.07


FIG. 12C

MHQ-86




8.22

MS-139




7.94

MHQ-94




8.04

MHQ-60




7.86

MS-91




7.84

MS-53




7.84

MS-96



7.81


MHQ-111



7.81


FIG. 12D

MS-86




7.80

MS-127




7.80

MS-48



7.77

MS-101



7.76

FIG. 12E

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/16723

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D215/44 C07D219/12 A61K31/47 A61P37/06

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	STREKOWSKI L ET AL: "Structure-activity relationship analysis of substituted 4-quinolinamines, antagonists of immunostimulatory CpG-oligodeoxynucleotides" BIOORGANIC & MEDICINAL CHEMISTRY LETTERS, GB, OXFORD, vol. 9, no. 13, 5 July 1999 (1999-07-05), pages 1819-1824, XP004168845 ISSN: 0960-894X the whole document	1,2,4, 6-8,11, 29,30, 32, 34-36,39
X	US 5 304 554 A (L. STREKOWSKI ET AL) 19 April 1994 (1994-04-19) column 8 -column 9; tables 4,5 -/--	1,4,29, 32

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

8 November 2000

Date of mailing of the international search report

24/11/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel: (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Siatou, E

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/16723

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 886 185 A (HAN-YEN CHOU ET AL) 23 March 1999 (1999-03-23) the whole document ---	49,50
A	D. E. MACFARLANE ET AL: "Antagonism of Immunostimulatory CpG-Oligodeoxynucleotides by quinacrine, Chloroquine, and Structurally Related Compounds" THE JOURNAL OF IMMUNOLOGY, vol. 160, 1998, pages 1122-1131, XP002152352 cited in the application page 1127; figure 9 page 1128; table II page 1130, left-hand column, line 27 -right-hand column, line 4 ---	1-14, 29-42
A	L. STREKOWSKI ET AL: "Synthesis and Structure-DNA Binding Relationship Analysis o DNA Triple-Helix Specific Intercalators" JOURNAL OF MEDICINAL CHEMISTRY, vol. 39, 1996, pages 3980-3983, XP002152353 cited in the application page 3981; table 1 ---	1-10, 29-33
A	EP 0 607 439 A (EISAI CO. LTD.) 27 July 1994 (1994-07-27) examples 256-259 page 254 claims 1,4 ---	1-14, 29-42
A	WO 93 07126 A (BOARD OF REGENTS OF THE UNIVERSITY OF NEBRASKA) 15 April 1993 (1993-04-15) claims 1-20 ---	43-48
A	WO 95 35287 A (F. HOFFMANN-LA ROCHE AG) 28 December 1995 (1995-12-28) claims 1-6,8-11 -----	43-48

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
PCT/US 00/16723

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5304554	A	19-04-1994	NONE	
US 5886185	A	23-03-1999	NONE	
EP 607439	A	27-07-1994	AU 668363 B	02-05-1996
			AU 2685192 A	03-05-1993
			FI 941417 A	25-03-1994
			KR 138695 B	01-10-1998
			NO 941101 A	30-05-1994
			US 5576322 A	19-11-1996
			CA 2116336 A	15-04-1993
			CN 1071164 A	21-04-1993
			HU 70854 A	28-11-1995
			WO 9307124 A	15-04-1993
			JP 10095776 A	14-04-1998
			JP 2818487 B	30-10-1998
			MX 9205542 A	31-03-1993
			NZ 244526 A	26-07-1995
			PT 100905 A	28-02-1994
			US 5693652 A	02-12-1997
			US 5801180 A	01-09-1998
			ZA 9207465 A	13-04-1993
WO 9307126	A	15-04-1993	US 5510356 A	23-04-1996
			AU 3176393 A	03-05-1993
WO 9535287	A	28-12-1995	AU 691306 B	14-05-1998
			AU 2736995 A	15-01-1996
			CA 2192203 A	28-12-1995
			EP 0765313 A	02-04-1997
			JP 10501540 T	10-02-1998
			NZ 288223 A	24-11-1997
			US 5736557 A	07-04-1998
			ZA 9504796 A	03-01-1996